

Current Topics in Microbiology and Immunology

Ben Adler *Editor*

# *Leptospira* and Leptospirosis

 Springer

# Current Topics in Microbiology and Immunology

Volume 387

## Series editors

Rafi Ahmed

School of Medicine, Rollins Research Center, Emory University, Room G211, 1510 Clifton Road, Atlanta, GA 30322, USA

Klaus Aktories

Medizinische Fakultät, Institut für Experimentelle und Klinische Pharmakologie und Toxikologie, Abt. I, Albert-Ludwigs-Universität Freiburg, Albertstr. 25, 79104 Freiburg, Germany

Richard W. Compans

Department of Microbiology and Immunology, Emory University, 1518 Clifton Road, CNR 5005, Atlanta, GA 30322, USA

Max D. Cooper

Department of Pathology and Laboratory Medicine, Georgia Research Alliance, Emory University, 1462 Clifton Road, Atlanta, GA 30322, USA

Jorge E. Galan

Boyer Ctr. for Molecular Medicine, School of Medicine, Yale University, 295 Congress Avenue, room 343, New Haven, CT 06536-0812, USA

Yuri Y. Gleba

ICON Genetics AG, Biozentrum Halle, Weinbergweg 22, 06120 Halle, Germany

Tasuku Honjo

Faculty of Medicine, Department of Medical Chemistry, Kyoto University, Sakyo-ku, Yoshida, Kyoto 606-8501, Japan

Yoshihiro Kawaoka

Influenza Research Institute, University of Wisconsin-Madison, 575 Science Drive, Madison, WI 53711, USA

Bernard Malissen

Centre d'Immunologie de Marseille-Luminy, Parc Scientifique de Luminy, Case 906, 13288, Marseille Cedex 9, France

Michael B. A. Oldstone

Department of Immunology and Microbial Science, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

Rino Rappuoli

Novartis Vaccines, Via Fiorentina 1, Siena 53100, Italy

Peter K. Vogt

Department of Molecular and Experimental Medicine, The Scripps Research Institute, 10550 North Torrey Pines Road, BCC-239, La Jolla, CA 92037, USA

Honorary Editor: Hilary Koprowski (deceased)

Formerly at Biotechnology Foundation, Inc., Ardmore, PA, USA

More information about this series at <http://www.springer.com/series/82>

Ben Adler  
Editor

# *Leptospira* and Leptospirosis

 Springer

*Editor*  
Ben Adler  
Australian Research Council Centre  
of Excellence in Structural and  
Functional Microbial Genomics  
Monash University  
Clayton, VIC  
Australia

ISSN 0070-217X                      ISSN 2196-9965 (electronic)  
ISBN 978-3-662-45058-1          ISBN 978-3-662-45059-8 (eBook)  
DOI 10.1007/978-3-662-45059-8

Library of Congress Control Number: 2014954598

Springer Heidelberg New York Dordrecht London

© Springer-Verlag Berlin Heidelberg 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media ([www.springer.com](http://www.springer.com))

# Preface

It is appropriate that this volume should appear on the 100th anniversary of the isolation of *Leptospira* and its discovery as the causative agent of Weil's disease. The last 10 years have seen a resurgence of research activity on *Leptospira*, probably as a result of the availability of whole genome sequences and the development of genetic tools for the manipulation of pathogenic leptospires. The previous decade has seen double the number of papers published on *Leptospira* or leptospirosis than in any previous decade and more than all the publications in the first 50 years of leptospirosis research. It is gratifying that this activity has been accompanied by an increased awareness of the serious disease caused in humans and animals by this global pathogen.

In the years since the publication of "*Leptospira* and Leptospirosis" by Faine et al. (1999, MediSci, Melbourne) it became apparent that it was unlikely that a single person, or even a small group of authors, would be able to find the time to write an updated version. However, although much of the information in that book remains relevant, large parts have become so out of date as to be misleading. The solution to both these problems was to assemble a group of world experts on leptospirosis to contribute to this present volume, which brings together just such a group. There is of necessity some overlap between chapters. This is unavoidable; for example, how can one write about the leptospiral outer membrane without discussing proteins and LPS, which are also key players in pathogenesis and in interactions with the host immune system? The overlap is also desirable, in that each chapter can be read on a stand-alone basis, with reference to other chapters where appropriate.

There are many people to whom I wish to express my heartfelt thanks. First and foremost, I am grateful to the chapter authors in this volume for the alacrity and enthusiasm with which they accepted the invitation to contribute. To my many colleagues and associates over the past decades (too numerous to detail here), I appreciate your willingness to collaborate and to share your wisdom and insight. Many of you are contributors to this volume. However, I would like to express my

particular gratitude to Solly Faine, who has been a mentor, colleague and friend for over 44 years, and who first introduced me to that fascinating organism, the leptospire. Finally, my grateful thanks to my wife Stephanie for her love, patience and forbearance during the preparation of this volume and over the last 40 years.

Melbourne, Australia, October 2014

Ben Adler

# Contents

<b>History of Leptospirosis and <i>Leptospira</i></b> . . . . .	1
Ben Adler	
<b>Systematics of <i>Leptospiraceae</i></b> . . . . .	11
Paul N. Levett	
<b>Leptospiral Structure, Physiology, and Metabolism</b> . . . . .	21
Caroline E. Cameron	
<b>Genomics, Proteomics, and Genetics of <i>Leptospira</i></b> . . . . .	43
Mathieu Picardeau	
<b>Leptospirosis in Humans</b> . . . . .	65
David A. Haake and Paul N. Levett	
<b>Animal Leptospirosis</b> . . . . .	99
William A. Ellis	
<b>The Molecular Basis of Leptospiral Pathogenesis</b> . . . . .	139
Gerald L. Murray	
<b>The Leptospiral Outer Membrane</b> . . . . .	187
David A. Haake and Wolfram R. Zückert	
<b>Host Response to <i>Leptospira</i> Infection</b> . . . . .	223
Richard L. Zuerner	
<b>Vaccines Against Leptospirosis</b> . . . . .	251
Ben Adler	



<b>The Role of Leptospirosis Reference Laboratories</b> . . . . .	273
Rudy A. Hartskeerl and Lee D. Smythe	
<b>Index</b> . . . . .	289

# History of Leptospirosis and *Leptospira*

Ben Adler

**Abstract** *Leptospira* was isolated and identified as the causative agent of the severe human syndrome Weil's disease about 100 years ago almost simultaneously, but independently, by workers in Japan and Europe. Since that time leptospires have been isolated from almost all mammalian species on every continent except Antarctica, with leptospirosis now recognized as the most widespread zoonosis worldwide and also a major cause of disease in many domestic animal species. Recent advances in molecular taxonomy have facilitated the development of a rational classification system, while the availability of genome sequences and the development of mutagenesis systems have begun to shed light on mechanisms of pathogenesis that appear to be unique to *Leptospira*.

## Contents

1 History of Weil's Disease.....	1
2 A Spirochete as the Causative Agent.....	3
3 Rats as Carriers of <i>Leptospira</i> .....	6
4 Recognition of Leptospirosis in Animals and the Expansion of Serovars and Syndromes.....	7
5 Nomenclature and Classification.....	7
6 Recent Developments.....	8
References.....	8

## 1 History of Weil's Disease

The modern history of leptospirosis began in 1886 when Adolph Weil (Fig. 1) described a particular type of jaundice accompanied by splenomegaly, renal dys-

---

B. Adler (✉)

Department of Microbiology, Australian Research Council Centre of Excellence in Structural and Functional Microbial Genomics, Monash University, Clayton, VIC 3800, Australia  
e-mail: Ben.Adler@monash.edu

© Springer-Verlag Berlin Heidelberg 2015

B. Adler (ed.), *Leptospira and Leptospirosis*, Current Topics in Microbiology and Immunology 387, DOI 10.1007/978-3-662-45059-8\_1

1

**Fig. 1** Portrait of Adolph Weil (1848–1916). Image courtesy of Wellcome Library, London



function, conjunctivitis, and skin rashes (Weil 1886). It was subsequently named Weil's disease. Although the etiology of the disease was unknown, it appeared to be infectious in nature and was often associated with outdoor occupations in which persons came into contact with water. Epidemics were common among sewer workers, rice-field workers, and coal miners.

However, it is apparent that leptospirosis had existed for millennia. Although it is difficult to draw firm conclusions from records before the advent of modern medical and scientific literature, it seems clear that at least some of the early disease outbreaks described in ancient texts were leptospirosis. For example, ancient Chinese texts carry accounts of "rice field jaundice", while in Japan syndromes clearly recognizable today as leptospirosis were termed "autumn fever" or "seven-day fever" (Kitamura and Hara 1918). In Europe, Australia and elsewhere, associations were recognized between febrile illness and particular occupations, giving rise to syndromes such as "cane-cutter's disease", "swine-herd's disease", and "Schlammfieber (mud fever)", well before the common etiology was recognized and identified (Alston and Broom 1958; van Thiel 1948). For a more detailed description of the early accounts of what were almost certainly large-scale outbreaks of leptospirosis, the reader is referred to Chapter 1 of Faine et al. (1999).


## 2 A Spirochete as the Causative Agent

Although *Leptospira* was first isolated independently and almost simultaneously in Japan and in Europe (see below), it is clear that the first demonstration of leptospire was made some years earlier by Stimson (1907), who used the recently described Levaditi silver deposition staining technique to observe spirochetes in kidney tissue sections of a patient described as having died of yellow fever (Figs. 2 and 3). It is probable that the patient was convalescing from Weil's disease when he contracted fatal yellow fever; spirochetes were observed in kidney, but not liver or heart, tissues. Stimson called the organism *Spirocheta interrogans*; the species name, which survives to this day, was suggested by the resemblance of the bacterial cells to a question mark, a feature that we now know to be due to the characteristic hooked ends of leptospires.

The first isolation of *Leptospira* followed just a few years later. In Japan, where Weil's disease was common in coal miners, Inada et al. (Fig. 4) injected guinea-pigs intraperitoneally with the blood of Weil's disease patients and succeeded in reproducing typical, acute leptospirosis in the animals (Inada et al. 1916). This and subsequent papers constituted a *tour de force* for the period; they defined transmissibility, routes of infection, pathological changes, tissue distribution, urinary excretion, leptospiral filterability, morphology, and motility. Signs in infected guinea-pigs included jaundice, conjunctivitis, inappetence, anemia, hemorrhages, and albuminuria. Disease was transferred in guinea-pigs for up to 50 generations. Spirochetes were observed in most tissues, with liver and kidneys containing the greatest numbers. These observations were extended to postmortem tissues from human cases, which revealed similar findings. These workers also showed that rabbits, mice, and rats were comparatively resistant to acute disease, even when injected with very large volumes of infected guinea-pig tissues.

Within a few months Inada and colleagues had succeeded in propagating the spirochetes in vitro in a medium made from emulsified guinea-pig kidney, and showed a preference for growth at 25 °C, with loss of viability at 37 °C.

**Fig. 2** Stimson's original observation of spirochetes in kidney tissue. Reproduced from (Noguchi 1928), with permission from the publishers, University of Chicago Press



## PUBLIC HEALTH REPORTS.

### UNITED STATES.

#### NOTE ON AN ORGANISM FOUND IN YELLOW-FEVER TISSUE.

[By Asst. Surg. A. M. Stimson, Hygienic Laboratory, Public Health and Marine-Hospital Service.]

The spirochætal origin of yellow fever has been suggested by Schaudinn and by Novy, but so far as I know no organism belonging to the genus *Spirochaeta* has been reported as having been found in yellow-fever tissues. I have examined some tissues for this class of organisms, using the method of Levaditi, as employed for the demonstration of *Treponema pallidum* in syphilis. The available yellow-fever material was unfortunately limited to the tissues from one case, all other specimens having been fixed otherwise than in formalin, which is required by Levaditi's method. The brain, liver, heart, and kidneys were examined. Nothing of special interest was observed in the first three, but in the kidney a very definite organism was found, having the following characters: Color, opaque black, in sharp contrast to the surrounding tissues; general appearance strongly suggesting a *spirochæte*; often irregularly curved, but some individuals having a regular series of alternate curves, no other indication of segments being observed; one or both extremities often bent back in the form of a hook; entire length variable up to  $14\ \mu$  or more; width estimated at  $\frac{1}{2}\ \mu$ ; length of short curves,  $1\frac{1}{2}$  to  $2\ \mu$ . It was confined to the cells and lumina of the tubules of the kidney, none being observed in the blood vessels, glomeruli, or interstitial tissue. While in some fields no organisms were present, in others large numbers were crowded together or scattered individuals were observed. The kidney from another case of yellow fever, where the tissue had been fixed in bichloride of mercury and acetic acid, was examined, and while no stained organisms were found, structures resembling those described were present in an unstained condition. The organisms were not found in kidney tissue from a patient dying of malaria, although the tissue was properly fixed and stained.

The object of this report is simply to invite attention to the findings in a single case, since no opinion as to the significance of this organism can be formed from such meager data. For convenience of reference I would suggest that the organism be called (*?Spirochaeta*) *interrogans*, the specific name being suggested by the form, somewhat resembling a question mark, which the organism frequently assumed in my preparations.

Fig. 3 Copy of Stimson's (1907) article in Public Health Reports. US Public Domain

The organism was named *Spirochaeta icterohaemorrhagiae*. One of the first isolates survives to this day and Ictero No. 1 was accepted by the Subcommittee on the Taxonomy of *Leptospira* in 1990 as the Type Strain of *Leptospira interrogans* (Marshall 1992).

**Fig. 4** Portrait of Ryokichi Inada (1874–1950). Kindly provided by Prof. Shin-ichi Yoshida, Kyushu University



Remarkably, the Japanese group also conducted the first vaccination studies. It is worth quoting verbatim.

Guinea pigs were immunized with repeated injections of liver emulsion of the infected animal and later with a pure culture of the spirochete, which had been killed by carbolic acid [Author's note, phenol]. The animals thus immunized did not develop the disease on the injection of the spirochete, which, it was known, would produce the disease in healthy animals. Hence this method seems promising for the prevention of the disease in man. Our conclusion is that the flea and mosquito have no share in the infection.

Of course, in the absence of quantitative data it is impossible to assess the degree of protection.

Finally, Inada and colleagues demonstrated immune lysis of leptospire by patient serum within the guinea-pig peritoneal cavity (the so-called Pfeiffer's method) and showed passive protection of guinea-pigs by convalescent patient serum or immune goat serum, but only if it was administered before the onset of jaundice. The importance of early treatment before the onset of organ failure remains relevant today (see the chapters by [D.A. Haake](#) and [P.N. Levett](#) and [W.A. Ellis](#), this volume).

The 1916 paper of Inada et al. extended data that were first published in the Japanese literature in early 1915 which described their observation in November 1914 of leptospire in the liver of a guinea-pig injected with the blood of a Weil's disease patient. Of course, this work was not known in Europe where trench warfare in World War I resulted in large numbers of Weil's disease cases. Two German groups independently and almost simultaneously (October 1915) succeeded in transmitting the infection to guinea-pigs and demonstrating the leptospire in guinea-pig tissues (Hubener 1915; Uhlenhuth and Fromme 1915). The groups named the organism *Spirochaeta nodosa* and *Spirochaeta icterogenes* respectively. Some controversy followed about priority, but it is clear that the Japanese discovery pre-dates the European ones by about a year, a fact recognized by the Subcommittee on the Taxonomy of *Leptospira* in specifying Ictero No. 1 as the type strain.

### 3 Rats as Carriers of *Leptospira*

The key finding that rats were renal carriers of *Leptospira* followed within 2 years, also reported by the Japanese group (Ido et al. 1917). The investigation was prompted by the serendipitous findings of spirochetes in the kidneys of field mice by colleagues working on tsutsugamushi (now *Orientia tsutsugamushi*). Ido and colleagues observed and cultured spirochetes from the kidneys and urine of a range of species of house and wild rats and identified them as *S. icterohaemorrhagiae* based on specific Pfeiffer reactivity with immune serum. They also made the key observation that leptospire were restricted to the kidneys and that the rats appeared healthy, the first observation of the asymptomatic carrier state. The connection between rats and Weil's disease was clearly established, as in coal mines which were frequently infested with rats, and also with the following epithet: "Cooks working in kitchens frequented by rats often became ill with spirochetosis icterohaemorrhagica." Interestingly, the group also observed spirochetes in mouse kidneys, but they were much less virulent when injected into guinea-pigs. It is probable that they observed one of the several serovars that we now know to be carried by mice. The Japanese findings were quickly confirmed in Europe and the U.S.A (Noguchi 1917; Stokes et al. 1917).

The Japanese group also reported some interesting epidemiological observations. Weil's disease in Japan showed a clear increased incidence in spring and autumn, but in coal mines where there was no temperature fluctuation the prevalence was the same year round. While this difference may be explained partly by season-specific human activities, the point was noted that higher incidence corresponded to temperatures of 22–25 °C. In addition, the incidence in coal mines with neutral or alkaline soil and water was high, whereas in mines with acidic soil and water infection was rare, despite equally high levels of rat infestation.