Human PHYSIOLOGY

Fourteenth Edition

Mc Graw Hill Education

Stuart Ira Fox

e me

me me !!

6A0

Human PHYSIOLOGY

Stuart Ira Fox Pierce College





HUMAN PHYSIOLOGY, FOURTEENTH EDITION

Published by McGraw-Hill Education, 2 Penn Plaza, New York, NY 10121. Copyright © 2016 by McGraw-Hill Education. All rights reserved. Printed in the United States of America. Previous editions © 2013, 2011, and 2009. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written consent of McGraw-Hill Education, including, but not limited to, in any network or other electronic storage or transmission, or broadcast for distance learning.

Some ancillaries, including electronic and print components, may not be available to customers outside the United States.

This book is printed on acid-free paper.

1 2 3 4 5 6 7 8 9 0 DOW/DOW 1 0 9 8 7 6 5

ISBN 978-0-07-783637-5 MHID 0-07-783637-5

Senior Vice President, Products & Markets: Kurt L. Strand Vice President, General Manager, Products & Markets: Marty Lange Vice President, Content Design & Delivery: Kimberly Meriwether David Managing Director: Michael Hackett Director of Digital Content: Michael G. Koot, PhD Brand Manager: Amy Reed/Chloe Bouxsein Director, Product Development: Rose Koos Production Developer: Fran Simon Marketing Manager: Jessica Cannavo Digital Product Analyst: John J. Theobald Director, Content Design & Delivery: Linda Avenarius Program Manager: Angela R. FitzPatrick Content Project Managers: April R. Southwood/Sherry L. Kane Buyer: Sandy Ludovissy Design: Matt Backhaus Content Licensing Specialist: John Leland Cover Image: Bill Westwood Compositor: Laserwords Private Limited Printer: R. R. Donnelley

All credits appearing on page or at the end of the book are considered to be an extension of the copyright page.

Library of Congress Cataloging-in-Publication Data

Fox, Stuart Ira.
Human physiology/Stuart Ira Fox, Pierce College.—Fourteenth edition. pages cm
Includes index.
ISBN 978-0-07-783637-5 (alk. paper)
1. Human physiology—Textbooks. I. Title.
QP34.5.F68 2016
612—dc23

2014044416

The Internet addresses listed in the text were accurate at the time of publication. The inclusion of a website does not indicate an endorsement by the authors or McGraw-Hill Education, and McGraw-Hill Education does not guarantee the accuracy of the information presented at these sites.

www.mhhe.com

Brief Contents

- **1** The Study of Body Function 1
- 2 Chemical Composition of the Body 24
- **3** Cell Structure and Genetic Control 50
- 4 Enzymes and Energy 88
- **5** Cell Respiration and Metabolism 106
- 6 Interactions Between Cells and the Extracellular Environment 130
- **7** The Nervous System 162
- 8 The Central Nervous System 206
- **9** The Autonomic Nervous System 243
- 10 Sensory Physiology 266
- **11** Endocrine Glands 316
- **12** Muscle 359

- **13** Blood, Heart, and Circulation 404
- 14 Cardiac Output, Blood Flow, and Blood Pressure 450
- **15** The Immune System 493
- **16** Respiratory Physiology 532
- 17 Physiology of the Kidneys 581
- **18** The Digestive System 619
- **19** Regulation of Metabolism 661
- **20** Reproduction 701

Appendix Answers to Objective Questions A-1

Glossary G-1

Credits C-1


Index I-1

About the Author

Stuart Ira Fox earned a Ph.D. in human physiology from the Department of Physiology, School of Medicine, at the University of Southern California, after earning degrees at the University of California at Los Angeles (UCLA); California State University, Los Angeles; and UC Santa Barbara. He has spent most of his professional life teaching at Los Angeles City College; California State University, Northridge; and Pierce College, where he has won numerous teaching awards, including several Golden Apples. Stuart has authored thirty-nine editions of seven textbooks, which are used worldwide and have been translated into several languages, and two novels. When not engaged in professional activities, he likes to hike, fly fish, and cross-country ski in the Eastern Sierra Nevada Mountains.

I wrote the first edition of Human Physiology to provide my students with a readable textbook to support the lecture material and help them understand physiology concepts they would need later in their health curricula and professions. This approach turned out to have wide appeal, which afforded me the opportunity to refine and update the text with each new edition. Writing new editions is a challenging educational experience, and an activity I find immensely enjoyable. Although changes have occurred in the scientific understanding and applications of physiological concepts, the students using this fourteenth edition have the same needs as those who used the first, and so my writing goals have remained the same. I am thankful for the privilege of being able to serve students and their instructors through these fourteen editions of Human Physiology.

-Stuart Ira Fox




To my wife, Ellen; and to Laura, Eric, Kayleigh, and Jacob Van Gilder; for all the important reasons.

Preface

The Cover

William B. Westwood's cover illustration of the eye and the structures and processes required for vision encompasses the study of physiology at multiple levels. The physiology of vision entails the biophysical processes of light becoming focused onto and interacting with photoreceptors, the molecular and cellular constituents of these receptors that enable



them to respond to light, and neural interactions needed for the brain to meaningfully interpret this stimulation.

Photoreceptors are located in the part of the eye and brain called the retina, which is a neural layer at the back of the eye. The front cover shows light entering the eye and becoming focused by the lens onto the retina. The outer segments of photoreceptors contain stacks of membranes, shown as purple at the bottom of the book's spine, which contain the photoreceptor pigment rhodopsin (the green structures within the membranes at the bottom left of the front cover).

The bottom middle of the front cover illustrates a plasma membrane of a photoreceptor neuron containing ion channels (pink). In the dark, these channels allow Na^+ ions (pink spheres) to enter the photoreceptor. Light induces a change in the rhodopsin that initiates a signaling pathway (not shown), which leads to the closing of these channels (shown by the bottom channel). This indirectly causes the photoreceptors to stimulate other neurons in the retina (bipolar cells, depicted in red near the bottom of the front cover), which then stimulate another layer of neurons (ganglion cells, depicted green at the bottom of the front cover.).

The axons (nerve fibers) of the ganglion cells gather together to form the optic nerves, which leave the eye to carry visual information to the brain, as shown on the back cover. The visual fields illustrated as blue and purple circles on the back cover stimulate different regions of the retina. Because many of the axons in the optic nerves cross to the opposite side, aspects of the right visual field are conveyed to the left cerebral cortex and vice versa, as illustrated by the blue and purple colors of the nerve tracts. Physiological processes continue within the brain, allowing it to create images that our mind interprets as the reality of the external world.

What Sets This Book Apart?

The study of human physiology provides the scientific foundation for the field of medicine and all other professions related to human health and physical performance. The scope of topics included in a human physiology course is therefore wideranging, yet each topic must be covered in sufficient detail to provide a firm basis for future expansion and application.

Human Physiology, fourteenth edition, is written for the undergraduate introductory human physiology course. Based on the author's extensive experience with teaching this course, the framework of the textbook is designed to provide basic biology and chemistry (chapters 2–5) before delving into more complex physiological processes. This approach is appreciated by both instructors and students; specific references in later chapters direct readers back to the foundational material as needed, presenting a self-contained study of human physiology.

In addition to not presupposing student's preparedness, this popular textbook is known for its clear and approachable writing style, detailed realistic art, and unsurpassed clinical information.

Acknowledgments

Reviewers

Patti Allen, Dixie State College Dani Behonick, Canada College Justin Brown, James Madison University Michael Burg, San Diego City College Julia Chang, Mount St. Mary's College Chalon Corey Cleland, James Madison University Linda Collins, University of Tennessee Chattanooga Maria Elena DeBellard, California State University-Northridge Andrew Flick, James Madison University James Hoffmann, Diablo Valley College Cynthia Kay-Nishiyama, California State University–Northridge Paul Kingston, San Diego City College Arnold Kondo, Citrus College Ann Maliszewski, Cuesta College Nancy Mann, Cuesta College Tim Maze, Lander University Vikki Mccleary, University of North Dakota Cheryl Neudauer, Minneapolis Community & Technical College Mark Paternostro, West Virginia University-Morgantown Erik Schweitzer, Santa Monica Community College Laura Steele, Ivy Tech Community College of Indiana-Fort Wayne R. Douglas Watson, University of Alabama at Birmingham Allison Wilson, Benedictine University

GUIDED TOUR

WHAT MAKES THIS TEXT A MARKET LEADER?

Clinical Applications—No Other Human Physiology Text Has More!

The framework of this textbook is based on integrating clinically germane information with knowledge of the body's physiological processes. Examples of this abound throughout the book. For example, in a clinical setting we record electrical activity from the body: this includes action potentials (chapter 7, section 7.2); EEG (chapter 8, section 8.2); and ECG (chapter 13, section 13.5). We also record mechanical force in muscle contractions (chapter 12, section 12.3). We note blood plasma measurements of many chemicals to assess internal body conditions. These include measurements of blood glucose (chapter 1, section 1.2) and the oral glucose tolerance test (chapter 19, section 19.4); and measurements of the blood cholesterol profile (chapter 13, section 13.7). These are just a few of many examples the author includes that focus on the connections between the study of physiology and our health industry.

NEW CLINICAL INVESTIGATIONS IN ALL CHAPTERS! **Clinical Investigation Chapter-Opening Clinical Investigations, Clues,** and Summaries are diagnostic case studies found in Sheryl, an active 78-year-old, suddenly became greatly each chapter. Clues are given throughout and the case is fatigued and disoriented while skiing. When she was finally resolved at the end of the chapter. brought to the hospital, blood tests revealed elevated levels of LDH. AST. ALT. and the MB isoform of CK. Some of the new terms and concepts you will encounter include: Clinical Investigation SUMMARY · Enzymes, isoenzymes, coenzymes, and cofactors • LDH, AST, ALT, and CK The sudden onset of Sheryl's great fatigue and disorintation is cause for concern and warranted immediate **CLUES Clinical Investigation** redical attention. Examination of table 4.1 with refernce to the disorders indicated by elevated levels of Sheryl's blood tests reveal elevated levels of CPK, LDH, K, LDH, AST, and ALT reveal that they share one pos-AST, and ALT. ble cause in common-myocardial infarction (heart What enzymes do these letters indicate, and what ttack). This possibility is reinforced by the laboratory diseases do elevated blood levels of these enzymes ests demonstrating that she had elevated levels of the suggest? K-MB isoenzyme, which is released by damaged heart


How might these test results relate to Sheryl's symptoms?

Clinical Investigations are enhanced with even more clinical assessments available on McGraw-Hill Connect[®]. These Clinical Investigations are written by the author and are specific to each chapter. They will offer the students great insight into that specific chapter. See additional chapter 4 Clinical Investigation on *Enzyme* Tests to Diagnose Diseases in the Connect site for this text.

exercise of skiina.

ells, rather than the CK-BB or CK-MM isoenzymes. A

ossible myocardial infarction could explain Sheryl's udden onset of symptom while performing the intense



ALL APPLICATION BOXES ARE NEW OR UPDATED!

Clinical Application Boxes are in-depth boxed essays that explore relevant topics of clinical interest and are placed at key points in the chapter to support the surrounding material. Subjects covered include pathologies, current research, pharmacology, and a variety of clinical diseases.

FITNESS APPLICATION

Metabolic syndrome is a combination of abnormal measurements-including central obesity (excess abdominal fat), hypertension (high blood pressure), insulin resistance (prediabetes), type 2 diabetes mellitus, high plasma triglycerides, and high LDL cholesterol-that greatly increase the risk of coronary heart disease, stroke, diabetes mellitus, and other conditions. The incidence of metabolic syndrome has increased alarmingly in recent years because of the increase in obesity. Eating excessive calories, particularly in the form of sugars (including high fructose corn syrup), stimulates insulin secretion. Insulin then promotes the uptake of blood glucose into adipose cells, where (through lipogenesis) it is converted into stored triglycerides (see figs. 5.12 and 5.13). Conversely, the lowering of insulin secretion, by diets that prevent the plasma glucose from rising sharply, promotes lipolysis (the breakdown of fat) and weight loss.

CLINICAL APPLICATION

When diseases damage tissues, some cells die and release their enzymes into the blood. The activity of these enzymes, reflecting their concentrations in the blood plasma, can be measured in a test tube by adding their specific substrates. Because an increase in certain enzymes in the blood can indicate damage to specific organs, such tests may aid the diagnosis of diseases. An increase in a man's blood levels of the acid, phosphatase, for example, may result from disease of the prostate (table 4.1).

Fitness Application Boxes are readings that explore physiological principles as applied to well-being, sports medicine, exercise physiology, and aging. They are also placed at relevant points in the text to highlight concepts just covered in the chapter.

Learning Outcomes are numbered for easy referencing in digital material!

Learning Outcome numbers are tied directly to Checkpoint numbers!

LEARNING OUTCOMES

After studying this section, you should be able to:

- **2.** Describe the aerobic cell respiration of glucose through the citric acid cycle.
- **3.** Describe the electron transport system and oxidative phosphorylation, explaining the role of oxygen in this process.

CHECKPOINT

- **2a.** Compare the fate of pyruvate in aerobic and anaerobic cell respiration.
- **2b.** Draw a simplified citric acid cycle and indicate the high-energy products.
- 3a. Explain how NADH and FADH₂ contribute to oxidative phosphorylation.
- **3b.** Explain how ATP is produced in oxidative phosphorylation.

GUIDED TOUR

WHAT MAKES THIS TEXT A MARKET LEADER?


Writing Style—Easygoing, Logical, and Concise


The words in *Human Physiology*, fourteenth edition, read as if the author is explaining concepts to you in a one-on-one conversation, pausing now and then to check and make sure you understand what he is saying. Each major section begins with a short overview of the information to follow. Numerous **comparisons** ("Unlike the life of an organism, which can be viewed as a linear progression from birth to death, the life of a cell follows a cyclical pattern"), **examples** ("A callus on the hand, for example, involves thickening of the skin by hyperplasia due to frequent abrasion"), **reminders** ("Recall that each member of a homologous pair came from a different parent"), and **analogies** ("In addition to this 'shuffling of the deck' of chromosomes ...") lend the author's style a comfortable grace that enables readers to easily flow from one topic to the next.

Exceptional Art—Designed from the Student's Point of View

What better way to support such unparalleled writing than with high-quality art? Large, bright illustrations demonstrate the physiological processes of the human body beautifully in a variety of ways.

> Stepped-out art clearly depicts various stages or movements with numbered explanations.











Labeled photos placed side by side with illustrations allow diagrammatic detail and realistic application.

Macro-to-micro art helps students put context around detailed concepts.





FOURTEENTH EDITION CHANGES

What's New?

Human Physiology, fourteenth edition, incorporates a number of new and recently modified physiological concepts. This may surprise people who are unfamiliar with the subject; indeed, the author sometimes is asked if the field really changes much from one edition to the next. It does; that's one of the reasons physiology is so much fun to study. Stuart has tried to impart this sense of excitement and fun in the book by indicating, in a manner appropriate for this level of student, where knowledge is new and where gaps in our knowledge remain.

The list that follows indicates only the larger areas of text and figure revisions and updates. It doesn't indicate instances where passages were rewritten to improve the clarity or accuracy of the existing material, or smaller changes made in response to information from recently published journals and from the reviewers of the previous edition.

GLOBAL CHANGES:

- Each Clinical Investigation in every chapter of the textbook is new.
- Each of the Clinical Investigation Clues, in every chapter, is new.
- The Clinical Investigation Summaries at the ends of all chapters are new.
- Every Clinical Application box, in each and every chapter, has been rewritten and updated.
- Every Fitness Application box, in each and every chapter, has been rewritten and updated.

MAJOR CHANGES IN CHAPTERS

These are specific changes made in the individual chapters in addition to the global changes described above.

Chapter 1: The Study of Body Function

- Discussions of exfoliative cytology and Pap smear added.
- Discussions of embryonic stem cells, totipotency, and pluripotency added.

Chapter 3: Cell Structure and Genetic Control

- New figures 3.3, 3.4, 3.7, 3.9*a*, and 3.18.
- Descriptions of microtubules and autophagosomes updated.
- Updated discussion of mitochondria, including hereditary mitochondrial diseases.
- Updated and expanded discussion of the agranular endoplasmic reticulum and drug tolerance.
- Updated and expanded discussion of genes, including new description of retrotransposons.
- Updated discussion of microRNA and new description of circular RNA.
- Updated discussion of the medical uses of RNA interference.
- Updated discussion of epigenetic regulation and its significance.

Chapter 5: Cell Respiration and Metabolism

- Updated description of the respiratory assemblies and their functions.
- New discussion of inherited mitochondrial diseases.
- Updated discussion of metabolic syndrome.
- Updated and expanded discussion of brown fat.

Chapter 6: Interactions Between Cells and the Extracellular Environment

- New figure 6.22*b*.
- Updated discussion of dialysis and hemodialysis.

Chapter 7: The Nervous System: Neurons and Synapses

- Updated and expanded discussions of microglia, axon regeneration, neurotrophins, astrocytes, and of microglia.
- Discussion of the structure and function of gap junctions updated and expanded.
- Figure 7.23 updated and revised.
- Explanation of synaptic vesicle docking and exocytosis updated and expanded.
- Expanded Table 7.4.
- New discussion of different subtypes of muscarinic ACh receptors.
- Updated and expanded discussion of dopamine receptors and new discussion of atypical antipsychotic drugs.
- Updated discussion of inhibitory neurotransmitters.
- Expanded discussion of endocannabinoid neurotransmitters.
- New discussion of hydrogen sulfide as a neurotransmitter.

Chapter 8: The Central Nervous System

- New photos in figures 8.9, 8.17, and 8.18.
- Updated and expanded discussion of CSF formation and circulation.
- Updated discussion of neurogenesis in the adult brain.
- Updated discussion of the origin of the electroencephalogram.
- New discussion of transient ischemic attack and stroke.
- Updated description of brain areas involved in memory storage.
- Updated and expanded discussion of Alzheimer's disease.
- Updated and expanded discussion of the molecular mechanisms involved in memory formation.
- Updated and expanded discussion of the roles of dendritic spines and neurogenesis in memory formation.
- Updated discussion of the regulation of circadian rhythms.
- Updated discussion of the role of the nucleus accumbens in the reward pathway.
- Updated discussion of orexin and new discussion of hypnotic drugs.

Chapter 9: The Autonomic Nervous System

New discussion of β₃-adrenergic receptors added.

Chapter 10: Sensory Physiology

- New figures 10.10 and 10.14*a*.
- Updated and expanded discussions of nociceptors, afferent fiber categories, and spinal cord lamina.
- Discussion of salty taste updated.

- - Updated and expanded discussion of olfactory processing.
 - Discussion of the structure and function of the cochlea updated and expanded.
 - New discussion of the role of microsaccades in vision.
 - New discussion of direction sensitive ganglion cells in vision.

Chapter 11: Endocrine Glands: Secretion and Action of Hormones

- New photos in figures 11.24 and 11.26.
- Updated and expanded discussion of the different drugs used to treat breast cancer.
- Updated and expanded discussion of insulin receptor structure and function.
- **Revised rendering of insulin receptor in figure 11.11.**
- Updated and expanded discussion of anterior pituitary cells and the hormones they produce.
- Updated and expanded discussion of stress and glucocorticoid effects.
- Updated discussions of calcitonin and the pancreatic islets.
- New discussion of adipokines and myokines.

Chapter 12: Muscle: Mechanisms of Contraction and Neural Control

- Expanded discussion of motor end plates and new explanation of end plate potential.
- New figure 12.9*a*.
- New discussion of the SERCA pumps in muscle contraction and relaxation.
- New discussion of muscle glycogen and exercise.
- Updated discussion of muscle metabolism of fat during exercise.
- New discussion of myokines and irisin.
- Updated and expanded discussion of satellite cells in muscle regeneration and sarcopenia.
- Updated and expanded discussion of calcium-induced calcium release in cardiac muscle.
- New discussion of calcium puffs and sparks in smooth muscle contraction.
- New discussion of myosin light-chain phosphatase in smooth muscle relaxation.

Chapter 13: Blood, Heart, and Circulation

- New discussion of the dietary need for iron in erythropoiesis.
- Updated discussions of hepcidin and the intrinsic clotting pathway.
- Updated discussion of the role of platelets in blood clotting and the use of warfarin to inhibit blood clotting.
- Updated and expanded discussion of the origin of the pacemaker potential.
- New discussion of sinoatrial conduction pathways and ectopic foci.
- Updated discussion of calcium pumping in the regulation of the heartbeat.
- New figure 13.31.
- Updated discussion of atherosclerosis.
- Updated discussion of myocardial infarction and diet.
- Updated and expanded discussion of blood tests to detect myocardial infarction.
- New discussion of interstitial fluid and the extracellular matrix.

Chapter 14: Cardiac Output, Blood Flow, and Blood Pressure


- New comparison of the pulmonary and systemic circulations.
- Updated discussion of the effects of sympathetic and parasympathetic nerves on the cardiac rate.
- Expanded discussion on the resting cardiac rate.
- New discussion of the Anrep effect.
- New discussion of neurovascular coupling and functional hyperemia.
- New goals for the treatment of hypertension discussed.
- Updated discussion of the mechanisms responsible for hypertension.
- Updated discussion of the role of dietary salt in hypertension.

Chapter 15: The Immune System

- Updated and expanded discussion of epithelial membranes and immunity.
- New discussion of NOD-like receptors and immunity.
- Updated and expanded discussion of opsonization and phagocytosis.
- Updated discussions of interferons and of secondary lymphoid organs.
- Updated discussion of the effects of mast cell cytokines in local inflammation.
- Updated discussion of the roles of resident macrophages and neutrophils in an inflammation.
- New figure 15.9.
- Updated discussions of helper and regulatory T cells and antigenpresenting cells.
- Updated discussion of MHC class-1 and class-2 molecules.
- Updated discussion of immune response to viral infections.
- Figures 15.15, 15.17, and 15.18 revised.
- Updated and expanded discussions of memory T cells and of adjuvants.
- New discussion of intravenous immunoglobulin.
- New discussion of humanized monoclonal antibodies and adoptive cell transfer.
- New discussion of natural killer T cells.
- Updated discussion of autoimmune and allergic reactions.
- Updated and expanded discussion of contact dermatitis.

Chapter 16: Respiratory Physiology

- Updated description of alveoli structure and function.
- New figures 16.3 and 16.5.
- Revised discussion of surfactant and respiratory distress syndrome.
- Updated and expanded discussion of the function of the diaphragm in ventilation.
- Updated discussions of asthma and of the pulmonary capillaries.
- Updated and expanded discussion of the mechanisms of ventilation/perfusion matching.
- Revised discussion of pulmonary hypertension and cor pulmonale.
- Updated and expanded discussion of the central regulation of breathing.



- Updated and expanded discussions of the carotid bodies and the central chemoreceptors.
- Updated discussion of the role of nitric oxide in acclimatization to high altitude.

Chapter 17: Physiology of the Kidneys

- Updated discussion of glomerular structure and function.
- New figure 17.9.
- Updated discussion of the renal tubule transport of sodium and chloride.
- Revised discussion of the countercurrent multiplier system.
- Updated discussion of urea transporters and aquaporin channels in the vasa recta.
- Updated discussion of countercurrent exchange in the renal medulla.
- Updated and expanded discussion of the role of urea in concentrating the urine.
- New discussion of arginine vasopressin as the antidiuretic hormone, and updated discussion of its secretion.
- Revised organization of the sections on renal plasma clearance.
- Updated discussion of renal tubule potassium secretion.
- Updated discussion of the roles of kidney-generated angiotensin II.
- New discussion of B-type natriuretic peptide.
- Updated discussion of ammonia produced by the renal tubules.

Chapter 18: The Digestive System

- Revised figure 18.7 and new fig. 18.11.
- Updated discussion of the lower esophageal sphincter.
- New discussion of parietal cells and potassium recycling.
- Updated discussion of Paneth cells and intestinal stem cells.
- Updated and expanded discussion of the enteric nervous system.
- Updated discussion of intestinal slow waves and action potentials.
- Updated and expanded discussion of the origin and function of the intestinal microbiota.
- Updated and expanded discussion of the antimicrobial properties of the intestinal mucosa.
- New discussion of the gut-associated lymphoid tissue.
- New discussions of Clostridium difficile infections and fecal microbiota transplantation.
- Updated discussions of liver fibrosis and cirrhosis.
- Updated and expanded discussion of transport processes in the pancreatic acini.
- New discussion of the function of somatostatin secreted by the D cells of the pancreatic islets.
- New discussion of incretins in the regulation of insulin secretion.
- Updated discussion of CCK in the regulation of pancreatic juice secretion.
- Updated discussion of secretin action.
- Updated discussions of fat transport and fatty acid uptake.

Chapter 19: Regulation of Metabolism

- New figures 19.17 and 19.20*a*.
- New discussion of hypothermia and hypothermic circulatory arrest.

- Updated discussion of the formation of the superoxide radical.
- Updated discussions of adipocyte turnover, and adipose tissue in starvation and obesity.
- Discussion of weight-loss medications updated.
- Updated and expanded discussion of hypothalamic neurons and neurotransmitters involved in the regulation of eating.
- Updated discussion of leptin and its regulation of appetite.
- New discussion of beige (or brite) adipocytes.
- Updated discussion of the mechanisms of beta cell insulin secretion.
- Updated discussion of how autonomic nerves and somatostatin regulate insulin secretion.
- Updated and expanded discussions of type 1 and type 2 diabetes and their treatments.
- New discussion on the roles of ectopic fat and visceral obesity in impaired glucose tolerance and type 2 diabetes.
- New discussion of soluble and insoluble fiber and its affect on insulin resistance.
- Updated discussion of dwarfism and new discussion of achondroplasia.
- Updated discussion of the regulation of osteoclast formation.
- New discussion of articular cartilage regeneration.
- Discussion of calcitonin updated.
- New discussion of osteocalcin and updated discussion of leptin actions on bone.
- Updated and expanded discussion of intestinal calcium absorption and the actions of vitamin D.
- Updated discussion of the actions of parathyroid hormone on renal phosphate excretion.


Chapter 20: Reproduction

- New figures 20.3, 20.40, and 20.42*c*.
- Updated discussion of X chromosome inactivation and SRY.
- New discussion of kisspeptins and the regulation of GnRH secretion.
- Updated discussion of DHT and estradiol in male physiology.
- Updated discussion of spermatogenesis and the blood-testis barrier.
- Updated and expanded discussions of the mechanisms of penile erection and of male contraception.
- Updated and expanded discussion of ovarian follicle hormone production and its regulation.
- Updated and expanded discussion of female contraception.
- Updated and expanded discussion of sperm capacitation and hyperactivation.
- New discussion of CatSper channels in sperm.
- Updated discussion of fertilization.
- Updated and expanded discussion of cloning and pluripotency.
- Updated discussion of stem cells in regenerative medicine.
- Updated discussion of adult stem cells and transdifferentiation.
- Updated and expanded discussion of the pituitary-like hormones secreted by the placenta.
- Table 20.7 updated and expanded.

Integrated and Adaptive Learning Systems



LearnSmartAdvantage.com



SMARTBOOK[®]

SmartBook is the first and only adaptive reading experience available for the higher education market. Powered by an intelligent diagnostic and adaptive engine, McGraw-Hill SmartBook facilitates the reading process by identifying what content a student knows and doesn't know through adaptive assessments. As the student reads, the reading material constantly adapts to ensure the student is focused on the content he or she needs the most to close any knowledge gaps.



LearnSmart is the only adaptive learning program proven to effectively assess a student's knowledge of basic course content and help him master it. By considering both confidence level and responses to actual content questions, McGraw-Hill LearnSmart identifies what an individual student knows and doesn't know and builds an optimal learning path, so that she spends less time on concepts she already knows and more time on those she doesn't. McGraw-Hill LearnSmart also predicts when a student will forget concepts and introduces remedial content to prevent this. The result is that McGraw-Hill Learn-Smart's adaptive learning path helps students learn faster, study more efficiently, and retain more knowledge, allowing instructors to focus valuable class time on higher-level concepts.




McGraw-Hill LearnSmart Labs is a superadaptive simulated lab experience that brings meaningful scientific exploration to students. Through a series of adaptive questions, McGraw-Hill LearnSmart Labs identifies a student's knowledge gaps and provides resources to quickly and efficiently close those gaps. Once students have mastered the necessary basic skills and concepts, they engage in a highly realistic simulated lab experience that allows for mistakes and the execution of the scientific method.



LEARNSMART

The primary goal of McGraw-Hill LearnSmart Prep is to help students who are unprepared to take college-level courses. Using super adaptive technology, the program identifies what a student doesn't know, and then provides "teachable moments" designed to mimic the office-hour experience. When combined with a personalized learning plan, an unprepared or struggling student has all the tools needed to quickly and effectively learn the foundational knowledge and skills necessary to be successful in a college-level course.



Continually evolving, McGraw-Hill Connect[®] has been redesigned to provide the only true adaptive learning experience delivered within a simple and easy-to-navigate environment, placing students at the very center.



Connect When is Connect

• Performance Analytics—Now available for both instructors and students, easy-to-decipher data illuminates course

performance. Students always know how they're doing in class, while instructors can view student and section performance at a glance.

- Mobile—Available on tablets. Students can now access assignments, quizzes, and results on the go, while instructors can assess student and section performance anytime, anywhere.
- Personalized Learning—Squeezing the most out of study time, the adaptive engine in Connect creates a highly personalized learning path for each student by identifying areas of weakness, and surfacing learning resources to assist in the moment of need. This seamless integration of reading, practice, and assessment ensures that the focus is on the most important content for that individual student at that specific time, while promoting long-term retention of the material.

http://connect.mheducation.com

Save time with auto-graded assessments and tutorials

Fully editable, customizable, auto-graded interactive assignments using high-quality art from the textbook, and animations and videos from a variety of sources take you way beyond multiple choice. Assignable content is available for every Learning Outcome in the book. Extremely high quality content, includes case study modules, animated learning modules, and more!

Generate powerful data related to student performance based on question tagging for Learning Outcomes, specific topics, Bloom's level, and more.

Everything you need, in one location

Enhanced Lecture Presentations contain lecture outlines, adjustable leader lines and labels, art, photos, tables, and animations embedded where appropriate. Fully customizable, but complete and ready to use, these presentations will enable you to spend less time preparing for lecture!


Animations—over 100 animations bringing key concepts to life, available for instructors and students.

Animation PPTs—animations are truly embedded in Power-Point[®] for ultimate ease of use! Just copy and paste into your custom slide show and you're done!

Tegrity R	ecorder			?
Hello, An	ny Reed		(Cha	ange User)
Course:	Select a course.			*
Title:	Week 2 Lecture	6		
Channel closed Channel closed Channel closed	ect Picture		Settings Recomma Audio Li	
Reco	rd a Class		Deg	rity

Take your course online—*easily*—with one-click Digital Lecture Capture

McGraw-Hill Tegrity[®] records and distributes your lecture with just a click of a button. Students can view them anytime/ anywhere via computer, tablet, or mobile device. Tegrity Campus indexes as it records your slideshow presentations and anything shown on your computer so students can use keywords to find exactly what they want to study.



Integrated and Adaptive Learning Systems



MH Campus[®] is an LMS integration service that offers instructors and students universal single signon, automatic registration, and gradebook synchronization with our learning platforms and content. Gain seamless access to our full library of digital assets—1,500 e-texts and instructor resources that let you build richer courses from within your chosen LMS!

Graw Hill Education Connectinsight

The first and only analytics tool of its kind, Connect Insight[®] is a series of visual data displays—each framed by an intuitive question—to provide at-a-glance information regarding how your class is doing.






Ph.I.L.S. 4.0 has been updated! Users have requested and we are providing *five new exercises* (Respiratory Quotient, Weight & Contraction, Insulin and Glucose Tolerance, Blood Typing, and Anti-Diuretic Hormone). **Ph.I.L.S. 4.0** is the perfect way to reinforce key physiology concepts with powerful Jab experiments. Created by Dr. Phil Stephens at Villanova University, this program

offers *42 laboratory simulations* that may be used to supplement or substitute for wet labs. All 42 labs are self-contained experiments—no lengthy instruction manual required. Users can adjust variables, view outcomes, make predictions, draw conclusions, and print lab reports. This easy-to-use software offers the flexibility to change the parameters of the lab experiment. There is no limit!

Fox's Laboratory Guide to Human Physiology: Concepts and Clinical Applications, 13th edition, contains time- and student-tested laboratory exercises that support most of the subjects covered in a human physiology course. It functions as a standalone manual, but is particularly well suited far use with Fox's Human Physiology, textbook.



Contents

Preface v

CHAPTER 1

The Study of Body Function 1

- 1.1Introduction to Physiology 2Scientific Method2
- **1.2 Homeostasis and Feedback Control 4** History of Physiology 4 Negative Feedback Loops 6 Positive Feedback 8 Neural and Endocrine Regulation 8 Feedback Control of Hormone Secretion 9
- **1.3The Primary Tissues10**Muscle Tissue11Nervous Tissue12Epithelial Tissue12Connective Tissue16
- **1.4 Organs and Systems 18** An Example of an Organ: The Skin 18
 Systems 20
 Body-Fluid Compartments 20

 Summary 21
 Review Activities 22

CHAPTER 2

Chemical Composition of the Body 24

- 2.1 Atoms, Ions, and Chemical Bonds 25 Atoms 25 Chemical Bonds, Molecules, and Ionic Compounds 26 Acids, Bases, and the pH Scale 29 Organic Molecules 30
- 2.2 Carbohydrates and Lipids 33 Carbohydrates 33 Lipids 36
- **2.3 Proteins 41** Structure of Proteins 41 Functions of Proteins 44

2.4 Nucleic Acids 44

Deoxyribonucleic Acid 44 Ribonucleic Acid 46 Summary 47 Review Activities 48

CHAPTER 3

Cell Structure and Genetic Control 50

- 3.1 Plasma Membrane and Associated Structures 51
 Structure of the Plasma Membrane 52
 Phagocytosis 54
 Endocytosis 54
 Exocytosis 55
 Cilia and Flagella 55
 Microvilli 56
 3.2 Cytoplasm and Its Organelles 56
 - Cytoplasm and Cytoskeleton 57 Lysosomes 58 Peroxisomes 58 Mitochondria 59 Ribosomes 60 Endoplasmic Reticulum 60 Golgi Complex 61

3.3 Cell Nucleus and Gene Expression 62 Genome and Proteome 63 Chromatin 63

RNA Synthesis 64 RNA Interference 67

3.4 Protein Synthesis and Secretion 68 Transfer RNA 68 Formation of a Polypeptide 69 Functions of the Endoplasmic Reticulum and Golgi Complex 70 Protein Degradation 70

3.5 DNA Synthesis and Cell Division 72 DNA Replication 72 The Cell Cycle 74 Mitosis 76 Meiosis 78 Epigenetic Inheritance 79

Interactions 83

Summary 84 Review Activities 85



Enzymes and Energy 88

4.1 Enzymes as Catalysts 89 Mechanism of Enzyme Action 89 Naming of Enzymes 91

4.2 Control of Enzyme Activity 92

 Effects of Temperature and pH 92
 Cofactors and Coenzymes 93
 Enzyme Activation 94
 Substrate Concentration and Reversible Reactions 94
 Metabolic Pathways 95

 4.3 Bioenergetics 97

Endergonic and Exergonic Reactions 98 Coupled Reactions: ATP 98 Coupled Reactions: Oxidation-Reduction 99 Summary 102 Review Activities 104

CHAPTER 5

Cell Respiration and Metabolism 106

5.1 Glycolysis and the Lactic Acid Pathway 107 Glycolysis 107 Lactic Acid Pathway 109

5.2 Aerobic Respiration 111

Citric Acid Cycle 111 Electron Transport and Oxidative Phosphorylation 112 Coupling of Electron Transport to ATP Production 113 ATP Balance Sheet 115

5.3 Interconversion of Glucose, Lactic Acid, and Glycogen 117

Glycogenesis and Glycogenolysis 117 Cori Cycle 117 5.4 Metabolism of Lipids and Proteins 119

Lipid Metabolism 119 Amino Acid Metabolism 122 Uses of Different Energy Sources 123

Interactions 126

Summary 127 Review Activities 128

CHAPTER 6

Interactions Between Cells and the Extracellular Environment 130

6.1 Extracellular Environment 131 Body Fluids 131 Extracellular Matrix 132 Categories of Transport Across the Plasma Membrane 132 6.2 Diffusion and Osmosis 133 Diffusion Through the Plasma Membrane 135 Rate of Diffusion 136 Osmosis 136 Regulation of Blood Osmolality 141 6.3 **Carrier-Mediated Transport** 142 Facilitated Diffusion 143 Active Transport 144 Bulk Transport 148 6.4 The Membrane Potential 149 Equilibrium Potentials 150 Resting Membrane Potential 152 6.5 Cell Signaling 153 Second Messengers 155 G-Proteins 155 Interactions 157 Summary 158 Review Activities 159

CHAPTER 7

The Nervous System 162

7.1 Neurons and Supporting Cells 163

Neurons 163 Classification of Neurons and Nerves 165 Neuroglial Cells 166 Neurilemma and Myelin Sheath 167 Functions of Astrocytes 170

- 7.2 Electrical Activity in Axons 172
 Ion Gating in Axons 173
 Action Potentials 174
 Conduction of Nerve Impulses 178
- 7.3 The Synapse 180Electrical Synapses: Gap Junctions 181Chemical Synapses 181
- 7.4 Acetylcholine as a Neurotransmitter 184 Chemically Regulated Channels 185 Acetylcholinesterase (AChE) 189 Acetylcholine in the PNS 190 Acetylcholine in the CNS 190
- 7.5 Monoamines as Neurotransmitters 191
 Serotonin as a Neurotransmitter 192
 Dopamine as a Neurotransmitter 192
 Norepinephrine as a Neurotransmitter 194
- 7.6 Other Neurotransmitters 194
 Amino Acids as Neurotransmitters 194
 Polypeptides as Neurotransmitters 196
 Endocannabinoids as Neurotransmitters 197
 Gases as Neurotransmitters 198
 ATP and Adenosine as Neurotransmitters 198
- **7.7 Synaptic Integration 199** Synaptic Plasticity 199 Synaptic Inhibition 200

Summary 201 Review Activities 203

CHAPTER 8

The Central Nervous System 206

- 8.1 Structural Organization of the Brain 207
- 8.2 Cerebrum 209
 - Cerebral Cortex 209 Basal Nuclei 215 Cerebral Lateralization 216 Language 218 Limbic System and Emotion 219 Memory 220 Emotion and Memory 224

8.3 Diencephalon 225

Thalamus and Epithalamus225Hypothalamus and Pituitary Gland226

8.4 Midbrain and Hindbrain 228 Midbrain 228 Hindbrain 230 Reticular Activating System in Sleep and Arousal 231

- 8.5 Spinal Cord Tracts 232 Ascending Tracts 233 Descending Tracts 233
- 8.6 Cranial and Spinal Nerves 236 Cranial Nerves 236 Spinal Nerves 236 Summary 239 Review Activities 240

CHAPTER 9

The Autonomic Nervous System 243

- 9.1 Neural Control of Involuntary Effectors 244 Autonomic Neurons 244 Visceral Effector Organs 245
 9.2 Divisions of the Autonomic Nervous
- 9.2 Divisions of the Autonomic Nervous System 246 Sympathetic Division 246

Parasympathetic Division 247

9.3 Functions of the Autonomic Nervous System 251

> Adrenergic and Cholinergic Synaptic Transmission 251 Responses to Adrenergic Stimulation 252 Responses to Cholinergic Stimulation 256 Other Autonomic Neurotransmitters 257 Organs with Dual Innervation 258 Organs Without Dual Innervation 260 Control of the Autonomic Nervous System by Higher Brain Centers 260

Interactions 262

Summary 263 Review Activities 264

CHAPTER 10

Sensory Physiology 266

10.1 Characteristics of Sensory Receptors 267 Categories of Sensory Receptors 267 Law of Specific Nerve Energies 268 Generator (Receptor) Potential 269 10.2 **Cutaneous Sensations** 270 Neural Pathways for Somatesthetic Sensations 271 Receptive Fields and Sensory Acuity 272 Lateral Inhibition 273 10.3 Taste and Smell 274 Taste 274 Smell 276 Vestibular Apparatus and Equilibrium 278 10.4 Sensory Hair Cells of the Vestibular Apparatus 279 Utricle and Saccule 280 Semicircular Canals 280 10.5 The Ears and Hearing 282 Outer Ear 283 Middle Ear 283 Cochlea 284 Spiral Organ (Organ of Corti) 286 10.6 The Eyes and Vision 290 Refraction 294 Accommodation 295 Visual Acuity 296 10.7 Retina 297 Effect of Light on the Rods 299 Electrical Activity of Retinal Cells 300 Cones and Color Vision 301 Visual Acuity and Sensitivity 304 Neural Pathways from the Retina 304 10.8 Neural Processing of Visual Information 307 Ganglion Cell Receptive Fields 307 Lateral Geniculate Nuclei 308 Cerebral Cortex 308 Interactions 310 Summary 311

Review Activities 314

CHAPTER **11**

Endocrine Glands 316

Endocrine Glands and Hormones 317 Chemical Classification of Hormones 318
 Prohormones and Prehormones 320
 Common Aspects of Neural and Endocrine
 Regulation 321
 Hormone Interactions 321

Effects of Hormone Concentrations on Tissue Response 321 11.2 Mechanisms of Hormone Action 323 Hormones That Bind to Nuclear Receptor Proteins 323 Hormones That Use Second Messengers 326 11.3 Pituitary Gland 331 Pituitary Hormones 331 Hypothalamic Control of the Posterior Pituitary 333 Hypothalamic Control of the Anterior Pituitary 333 Feedback Control of the Anterior Pituitary 335 Higher Brain Function and Pituitary Secretion 336 11.4 Adrenal Glands 337 Functions of the Adrenal Cortex 337 Functions of the Adrenal Medulla 339 Stress and the Adrenal Gland 340 11.5 Thyroid and Parathyroid Glands 341 Production and Action of Thyroid Hormones 342 Parathyroid Glands 344 11.6 Pancreas and Other Endocrine Glands 345 Pancreatic Islets (Islets of Langerhans) 345 Pineal Gland 346 Gastrointestinal Tract 349 Gonads and Placenta 349 11.7 Paracrine and Autocrine Regulation 349 Examples of Paracrine and Autocrine Regulation 350 Prostaglandins 351 Interactions 354

Summary 355 Review Activities 356

CHAPTER 12

Muscle 359

- 12.1 Skeletal Muscles 360 Structure of Skeletal Muscles 360 Motor End Plates and Motor Units 361
- **12.2 Mechanisms of Contraction 364** Sliding Filament Theory of Contraction 367 Regulation of Contraction 369
- **12.3** Contractions of Skeletal Muscles374Twitch, Summation, and Tetanus374

Types of Muscle Contractions 375 Series-Elastic Component 376 Length-Tension Relationship 376 12.4 **Energy Requirements of Skeletal** Muscles 377 Metabolism of Skeletal Muscles 378 Slow- and Fast-Twitch Fibers 380 Muscle Fatigue 381 Adaptations of Muscles to Exercise Training 382 Muscle Damage and Repair 384 12.5 Neural Control of Skeletal Muscles 384 Muscle Spindle Apparatus 386 Alpha and Gamma Motoneurons 387 Coactivation of Alpha and Gamma Motoneurons 387 Skeletal Muscle Reflexes 387 Upper Motor Neuron Control of Skeletal Muscles 390 12.6 Cardiac and Smooth Muscles 391 Cardiac Muscle 392 Smooth Muscle 393 Interactions 398

Summary 399 Review Activities 401

CHAPTER 13

Blood, Heart, and Circulation 404

13.1 Functions and Components of the Circulatory System 405

Functions of the Circulatory System405Major Components of the Circulatory System405

13.2 Composition of the Blood 406 Plasma 406

The Formed Elements of Blood 407 Hematopoiesis 409 Red Blood Cell Antigens and Blood Typing 412 Blood Clotting 414 Dissolution of Clots 417

13.3 Structure of the Heart 418 Pulmonary and Systemic Circulations 418 Atrioventricular and Semilunar Valves 419 Heart Sounds 420

13.4 Cardiac Cycle 422 Pressure Changes During the Cardiac Cycle 423 13.5 Electrical Activity of the Heart and the Electrocardiogram 425

> Electrical Activity of the Heart 425 The Electrocardiogram 428

13.6 Blood Vessels 431 Arteries 431 Capillaries 433

Veins 435

13.7 Atherosclerosis and Cardiac Arrhythmias 436

Atherosclerosis 436 Arrhythmias Detected by the Electrocardiograph 440

13.8 Lymphatic System 442

Summary 445 Review Activities 447

CHAPTER 14

Cardiac Output, Blood Flow, and Blood Pressure 450

14.1 Cardiac Output 451

Regulation of Cardiac Rate 451 Regulation of Stroke Volume 452 Venous Return 454

14.2 Blood Volume 456

Exchange of Fluid Between Capillaries and Tissues 457 Regulation of Blood Volume by the

Kidneys 459

14.3 Vascular Resistance to Blood Flow 463

Physical Laws Describing Blood Flow464Extrinsic Regulation of Blood Flow465Paracrine Regulation of Blood Flow466Intrinsic Regulation of Blood Flow467

14.4 Blood Flow to the Heart and Skeletal Muscles 468

Aerobic Requirements of the Heart 468 Regulation of Coronary Blood Flow 469 Regulation of Blood Flow Through Skeletal Muscles 470

Circulatory Changes During Exercise 470

14.5 Blood Flow to the Brain and Skin 473Cerebral Circulation 473Cutaneous Blood Flow 474

14.6 **Blood Pressure 475**

Baroreceptor Reflex 477 Atrial Stretch Reflexes 479 Measurement of Blood Pressure 479 Pulse Pressure and Mean Arterial Pressure 481

14.7 Hypertension, Shock, and Congestive Heart Failure 482 Hypertension 482

> Circulatory Shock 484 Congestive Heart Failure 486

Interactions 488

Summary 489 Review Activities 490

CHAPTER 15

The Immune System 493

15.1 **Defense Mechanisms** 494 Innate (Nonspecific) Immunity 495 Adaptive (Specific) Immunity 497 Lymphocytes and Lymphoid Organs 499 Local Inflammation 500

- 15.2 Functions of B Lymphocytes 503 Antibodies 504 The Complement System 506
- 15.3 Functions of T Lymphocytes 507 Killer, Helper, and Regulatory T Lymphocytes 507 Interactions Between Antigen-Presenting Cells and T Lymphocytes 511

Active and Passive Immunity 514 15.4

Active Immunity and the Clonal Selection Theory 515 Immunological Tolerance 517 Passive Immunity 518

- 15.5 Tumor Immunology 519 Natural Killer Cells 520 Effects of Aging and Stress 521
- 15.6 **Diseases Caused by the Immune** System 521 Autoimmunity 521

Immune Complex Diseases 522 Allerav 523

Interactions 527

Summarv 528 Review Activities 529

CHAPTER 16

Respiratory Physiology 532

16.1 The Respiratory System 533 Structure of the Respiratory System 533 Thoracic Cavity 536 **Physical Aspects of Ventilation 536** 16.2 Intrapulmonary and Intrapleural Pressures 537 Physical Properties of the Lungs 538 Surfactant and Respiratory Distress Syndrome 540 16.3 Mechanics of Breathing 540 Inspiration and Expiration 541 Pulmonary Function Tests 542 Pulmonary Disorders 544 16.4 Gas Exchange in the Lungs 547 Calculation of Po. 547 Partial Pressures of Gases in Blood 548 Significance of Blood Po, and Pco. Measurements 550 Pulmonary Circulation and Ventilation/Perfusion Ratios 550 Disorders Caused by High Partial Pressures of Gases 552 16.5 Regulation of Breathing 553

Brain Stem Respiratory Centers 553 Effects of Blood Pco. and pH on Ventilation 555 Effects of Blood Po, on Ventilation 557 Effects of Pulmonary Receptors on Ventilation 558

16.6 Hemoglobin and Oxygen Transport 559 Hemoglobin 559

The Oxyhemoglobin Dissociation Curve 561 Effect of pH and Temperature on Oxygen Transport 562

Effect of 2,3-DPG on Oxygen Transport 563 Inherited Defects in Hemoglobin Structure and Function 564

Muscle Myoglobin 564

- 16.7 Carbon Dioxide Transport 565 The Chloride Shift 566 The Reverse Chloride Shift 566
- 16.8 Acid-Base Balance of the Blood 567 Principles of Acid-Base Balance 568 Ventilation and Acid-Base Balance 569

16.9 Effect of Exercise and High Altitude on Respiratory Function 570

Ventilation During Exercise 570 Acclimatization to High Altitude 571

Interactions 575

Summary 576 Review Activities 578

CHAPTER 17

Physiology of the Kidneys 581

- 17.1 Structure and Function of the Kidneys 582
 Gross Structure of the Urinary System 582
 Control of Micturition 584
 Microscopic Structure of the Kidney 585
- **17.2 Glomerular Filtration 587** Glomerular Ultrafiltrate 588 Regulation of Glomerular Filtration Rate 588

17.3 Reabsorption of Salt and Water 590 Reabsorption in the Proximal Tubule 590 The Countercurrent Multiplier System 592 Collecting Duct: Effect of Antidiuretic Hormone (ADH) 595

17.4 Renal Plasma Clearance 598 Transport Process Affecting Renal

Clearance 599 Renal Clearance of Inulin: Measurement of GFR 600 Renal Clearance Measurements 601

Reabsorption of Glucose 603

17.5 Renal Control of Electrolyte and Acid-Base Balance 604

Role of Aldosterone in Na⁺/K⁺ Balance 604 Control of Aldosterone Secretion 606 Juxtaglomerular Apparatus 606 Natriuretic Peptides 607 Relationship Between Na⁺, K⁺, and H⁺ 608 Renal Acid-Base Regulation 608

17.6 Diuretics and Renal Function Tests 611 Use of Diuretics 611 Renal Function Tests and Kidney Disease 613

Interactions 614

Summary 615

Review Activities 616

CHAPTER 18

The Digestive System 619

- **18.1** Introduction to the Digestive System 620Layers of the Gastrointestinal Tract621Regulation of the Gastrointestinal Tract622
- 18.2 From Mouth to Stomach 623
 Esophagus 624
 Stomach 625
 Pepsin and Hydrochloric Acid Secretion 626
- **18.3 Small Intestine 628** Villi and Microvilli 629
 Intestinal Enzymes 630
 Intestinal Contractions and Motility 631

18.4 Large Intestine 632

Intestinal Microbiota 633 Fluid and Electrolyte Absorption in the Intestine 635 Defecation 636

18.5 Liver, Gallbladder, and Pancreas 636 Structure of the Liver 636 Functions of the Liver 638 Gallbladder 641 Pancreas 643

18.6 Regulation of the Digestive System 645

 Regulation of Gastric Function 645
 Regulation of Intestinal Function 648
 Regulation of Pancreatic Juice and Bile
 Secretion 648
 Trophic Effects of Gastrointestinal Hormones 650

18.7 Digestion and Absorption of Food 650

Digestion and Absorption of Carbohydrates650Digestion and Absorption of Proteins651Digestion and Absorption of Lipids652

Interactions 656

Summary 657 Review Activities 658

CHAPTER 19

Regulation of Metabolism 661

19.1 Nutritional Requirements 662 Metabolic Rate and Caloric Requirements 662 Anabolic Requirements 664

Vitamins and Minerals 666 Free Radicals and Antioxidants 668 19.2 **Regulation of Energy Metabolism 669** Regulatory Functions of Adipose Tissue 670 Regulation of Hunger and Metabolic Rate 672 Caloric Expenditures 674 Hormonal Regulation of Metabolism 675 19.3 **Energy Regulation by the Pancreatic** Islets 677 Regulation of Insulin and Glucagon Secretion 677 Insulin and Glucagon: Absorptive State 679 Insulin and Glucagon: Postabsorptive State 679 19.4 **Diabetes Mellitus and Hypoglycemia** 681 Type 1 Diabetes Mellitus 681 Type 2 Diabetes Mellitus 682 Hypoglycemia 685 19.5 Metabolic Regulation by Adrenal Hormones, Thyroxine, and Growth Hormone 686 Adrenal Hormones 686 Thyroxine 686 Growth Hormone 687 19.6 **Regulation of Calcium and Phosphate** Balance 690 Bone Deposition and Resorption 690 Hormonal Regulation of Bone 692 1,25-Dihydroxyvitamin D₃ 693 Negative Feedback Control of Calcium and Phosphate Balance 695 Summary 696 Review Activities 698



Reproduction 701

20.1 Sexual Reproduction 702

Sex Determination 702 Development of Accessory Sex Organs and External Genitalia 705 Disorders of Embryonic Sexual Development 706

20.2 Endocrine Regulation of Reproduction 708 Interactions Between the Hypothalamus, Pituitary Gland, and Gonads 709 Onset of Puberty 710

Human Sexual Response 712 20.3 Male Reproductive System 712 Control of Gonadotropin Secretion 713 Endocrine Functions of the Testes 714 Spermatogenesis 715 Male Accessory Sex Organs 718 Erection, Emission, and Ejaculation 719 Male Fertility 721 20.4 Female Reproductive System 722 **Ovarian Cycle 724 Ovulation 725** Pituitary-Ovarian Axis 727 **Menstrual Cycle 728** 20.5 Phases of the Menstrual Cycle: Cyclic Changes in the Ovaries 728 Cyclic Changes in the Endometrium 731 Effects of Pheromones, Stress, and Body Fat 732 Contraceptive Methods 733 Menopause 734 20.6 Fertilization, Pregnancy, and Parturition 734 Fertilization 735 Cleavage and Blastocyst Formation 737 Implantation of the Blastocyst and Formation of the Placenta 740 Exchange of Molecules Across the Placenta 742 Endocrine Functions of the Placenta 743 Labor and Parturition 744 Lactation 746 **Interactions 749**

Pineal Gland 712

Concluding Remarks 750

Summary 750 Review Activities 752

Appendix

Answers to Test Your Knowledge Questions A-1

Glossary G-1

Credits C-1

Index I-1

CHAPTER OUTLINE

- 1.1 Introduction to Physiology 2 Scientific Method 2
- **1.2 Homeostasis and Feedback Control 4** History of Physiology 4 Negative Feedback Loops 6 Positive Feedback 8 Neural and Endocrine Regulation 8 Feedback Control of Hormone Secretion 9

1.3 The Primary Tissues 10

Muscle Tissue 11 Nervous Tissue 12 Epithelial Tissue 12 Connective Tissue 16

1.4 Organs and Systems 18

An Example of an Organ: The Skin 18 Systems 20 Body-Fluid Compartments 20

Summary 21

Review Activities 22

CHAPTER

The Study of Body Function

Clinical Investigation

As you study the sections of chapter 1, you can see how your new knowledge can be applied to interesting health issues that may be important to know in your future career as a health professional. This can add zest to your studies and increase your motivation to truly understand physiological concepts, rather than to simply memorize facts for examinations. Each chapter begins with a medical mystery for you to solve, using information in the text of that chapter and "Clinical Investigation Clues" within the chapter.

For example, suppose Linda goes for a medical examination where her body temperature is measured, and she gives a fasting blood sample to test for glucose. Your first Clinical Investigation challenge is to determine the medical significance of these physiological tests.

1.1 INTRODUCTION TO PHYSIOLOGY

Human physiology is the study of how the human body functions, with emphasis on specific cause-and-effect mechanisms. Knowledge of these mechanisms has been obtained experimentally through applications of the scientific method.

LEARNING OUTCOMES

After studying this section, you should be able to:

- 1. Describe the scientific study of human physiology.
- 2. Describe the characteristics of the scientific method.

Physiology (from the Greek *physis* = nature; *logos* = study) is the study of biological function—of how the body works, from molecular mechanisms within cells to the actions of tissues, organs, and systems, and how the organism as a whole accomplishes particular tasks essential for life. In the study of physiology, the emphasis is on mechanisms—with questions that begin with the word *how* and answers that involve cause-and-effect sequences. These sequences can be woven into larger and larger stories that include descriptions of the structures involved (anatomy) and that overlap with the sciences of chemistry and physics.

The separate facts and relationships of these cause-andeffect sequences are derived empirically from experimental evidence. Explanations that seem logical are not necessarily true; they are only as valid as the data on which they are based, and they can change as new techniques are developed and further experiments are performed. The ultimate objective of physiological research is to understand the normal functioning of cells, organs, and systems. A related science—*pathophysiology*—is concerned with how physiological processes are altered in disease or injury.

Pathophysiology and the study of normal physiology complement one another. For example, a standard technique for investigating the functioning of an organ is to observe what happens when the organ is surgically removed from an experimental animal or when its function is altered in a specific way. This study is often aided by "experiments of nature" diseases—that involve specific damage to the functioning of an organ. The study of disease processes has thus aided our understanding of normal functioning, and the study of normal physiology has provided much of the scientific basis of modern medicine. This relationship is recognized by the Nobel Prize committee, whose members award prizes in the category "Physiology or Medicine."

The physiology of invertebrates and of different vertebrate groups is studied in the science of *comparative physiology*. Much of the knowledge gained from comparative physiology has benefited the study of human physiology. This is because animals, including humans, are more alike than they are different. This is especially true when comparing humans with other mammals. The small differences in physiology between humans and other mammals can be of crucial importance in the development of pharmaceutical drugs (discussed later in this section), but these differences are relatively slight in the overall study of physiology.

Scientific Method

All of the information in this text has been gained by people applying the scientific method. Although many different techniques are involved when people apply the scientific method, all share three attributes: (1) confidence that the natural world, including ourselves, is ultimately explainable in terms we can understand; (2) descriptions and explanations of the natural world that are honestly based on observations and that could be modified or refuted by other observations; and (3) humility, or the willingness to accept the fact that we could be wrong. If further study should yield conclusions that refuted all or part of an idea, the idea would have to be modified accordingly. In short, the scientific method is based on a confidence in our rational ability, honesty, and humility. Practicing scientists may not always display these attributes, but the validity of the large body of scientific knowledge that has been accumulated-as shown by the technological applications and the predictive value of scientific hypotheses-are ample testimony to the fact that the scientific method works.

The scientific method involves specific steps. After certain observations regarding the natural world are made, a **hypothesis** is formulated. In order for this hypothesis to be scientific, it must be capable of being refuted by experiments or other observations of the natural world. For example, one might hypothesize that people who exercise regularly have a lower resting pulse rate than other people. Experiments are conducted, or other observations are made, and the results are analyzed. Conclusions are then drawn as to whether the new data either refute or support the hypothesis. If the hypothesis survives such testing, it might be incorporated into a more general **theory.** Scientific theories are thus not simply conjectures; they are statements about the natural world that incorporate a number of proven hypotheses. They serve as a logical framework by which these hypotheses can be interrelated and provide the basis for predictions that may as yet be untested.

The hypothesis in the preceding example is scientific because it is *testable;* the pulse rates of 100 athletes and 100 sedentary people could be measured, for example, to see if there were statistically significant differences. If there were, the statement that athletes, on the average, have lower resting pulse rates than other people would be justified *based on these data*. One must still be open to the fact that this conclusion could be wrong. Before the discovery could become generally accepted as fact, other scientists would have to consistently replicate the results. Scientific theories are based on *reproduc-ible* data.

It is quite possible that when others attempt to replicate the experiment, their results will be slightly different. They may then construct scientific hypotheses that the differences in resting pulse rate also depend on other factors, such as the nature of the exercise performed. When scientists attempt to test these hypotheses, they will likely encounter new problems requiring new explanatory hypotheses, which then must be tested by additional experiments.

In this way, a large body of highly specialized information is gradually accumulated, and a more generalized explanation (a scientific theory) can be formulated. This explanation will almost always be different from preconceived notions. People who follow the scientific method will then appropriately modify their concepts, realizing that their new ideas will probably have to be changed again in the future as additional experiments are performed.

Use of Measurements, Controls, and Statistics

Suppose you wanted to test the hypothesis that a regular exercise program causes people to have a lower resting heart rate. First, you would have to decide on the nature of the exercise program. Then, you would have to decide how the heart rate (or pulse rate) would be measured. This is a typical problem in physiology research because the testing of most physiological hypotheses requires quantitative **measurements.**

The group that is subject to the testing condition—in this case, exercise—is called the **experimental group.** A measurement of the heart rate for this group would be meaningful only if it is compared to that of another group, known as the **control group.** How shall this control group be chosen? Perhaps the subjects could serve as their own controls—that is, a person's resting heart rate could be measured before and after the exercise regimen. If this isn't possible, a control group could be other people who do not follow the exercise program. The choice of control groups is often a controversial aspect of

physiology studies. In this example, did the people in the control group really refrain from *any* exercise? Were they comparable to the people in the experimental group with regard to age, sex, ethnicity, body weight, health status, and so on? You can see how difficult it could be in practice to get a control group that could satisfy any potential criticism.

Another possible criticism could be bias in the way that the scientists perform the measurements. This bias could be completely unintentional; scientists are human, after all, and they may have invested months or years in this project. To prevent such bias, the person doing the measurements often does not know if a subject is part of the experimental or the control group. This is known as a *blind measurement*.

Now suppose the data are in and it looks like the experimental group indeed has a lower average resting heart rate than the control group. But there is overlap—some people in the control group have measurements that are lower than some people in the experimental group. Is the difference in the average measurements of the groups due to a real physiological difference, or is it due to chance variations in the measurements? Scientists attempt to test the *null hypothesis* (the hypothesis that the difference is due to chance) by employing the mathematical tools of **statistics.** If the statistical results so warrant, the null hypothesis can be rejected and the experimental hypothesis can be deemed to be supported by this study.

The statistical test chosen will depend upon the design of the experiment, and it can also be a source of contention among scientists in evaluating the validity of the results. Because of the nature of the scientific method, "proof" in science is always provisional. Some other researchers, employing the scientific method in a different way (with different measuring techniques, experimental procedures, choice of control groups, statistical tests, and so on), may later obtain different results. The scientific method is thus an ongoing enterprise.

The results of the scientific enterprise are written up as research articles, and these must be reviewed by other scientists who work in the same field before they can be published in **peer-reviewed journals.** More often than not, the reviewers will suggest that certain changes be made in the articles before they can be accepted for publication.

Examples of such peer-reviewed journals that publish articles in many scientific fields include *Science* (www. sciencemag .org/), *Nature* (www.nature.com/nature/), and *Proceedings of the National Academy of Sciences* (www.pnas.org/). Review articles on physiology can be found in *Annual Review of Physiology* (physiol .annualreviews.org/), *Physiological Reviews* (physrev.physiology .org/), and *Physiology* (physiologyonline. physiology.org). Medical research journals, such as the *New England Journal of Medicine* (content.nejm.org/) and *Nature Medicine* (www.nature.com/nm/), also publish articles of physiological interest. There are also many specialty journals in areas of physiology such as neurophysiology, endocrinology, and cardiovascular physiology.

Students who wish to look online for scientific articles published in peer-reviewed journals that relate to a particular subject can do so at the National Library of Medicine website, *PubMed* (www.ncbi.nlm.nih.gov/entrez/query.fcgi).

Development of Pharmaceutical Drugs

The development of new pharmaceutical drugs can serve as an example of how the scientific method is used in physiology and its health applications. The process usually starts with basic physiological research, often at cellular and molecular levels. Perhaps a new family of drugs is developed using cells in tissue culture (*in vitro*, or outside the body). For example, cell physiologists studying membrane transport may discover that a particular family of compounds blocks membrane channels for calcium ions (Ca²⁺). Because of their knowledge of physiology, other scientists may predict that a drug of this nature might be useful in the treatment of hypertension (high blood pressure). This drug may then be tried in animal experiments.

If a drug is effective at extremely low concentrations *in vitro* (in cells cultured outside of the body), there is a chance that it may work *in vivo* (in the body) at concentrations low enough not to be toxic (poisonous). This possibility must be thoroughly tested utilizing experimental animals, primarily rats and mice. More than 90% of drugs tested in experimental animals are too toxic for further development. Only in those rare cases when the toxicity is low enough may development progress to human/clinical trials.

Biomedical research is often aided by **animal models** of particular diseases. These are strains of laboratory rats and mice that are genetically susceptible to particular diseases that resemble human diseases. Research utilizing laboratory animals typically takes several years and always precedes human (clinical) trials of promising drugs. It should be noted that this length of time does not include all of the years of "basic" physiological research (involving laboratory animals) that provided the scientific foundation for the specific medical application.

In phase I clinical trials, the drug is tested on healthy human volunteers. This is done to test its toxicity in humans and to study how the drug is "handled" by the body: how it is metabolized, how rapidly it is removed from the blood by the liver and kidneys, how it can be most effectively administered, and so on. If significant toxic effects are not observed, the drug can proceed to the next stage. In phase II clinical trials, the drug is tested on the target human population (for example, those with hypertension). Only in those exceptional cases where the drug seems to be effective but has minimal toxicity does testing move to the next phase. Phase III trials occur in many research centers across the country to maximize the number of test participants. At this point, the test population must include a sufficient number of subjects of both sexes, as well as people of different ethnic groups. In addition, people are tested who have other health problems besides the one that the drug is intended to benefit. For example, those who have diabetes in addition to hypertension would be included in this phase. If the drug passes phase III trials, it goes to the Food and

Drug Administration (FDA) for approval. **Phase IV trials** test other potential uses of the drug.

Less than 10% of the tested drugs make it all the way through clinical trials to eventually become approved and marketed. This low success rate does not count those that fail after approval because of unexpected toxicity, nor does it take into account the great amount of drugs that fail earlier in research before clinical trials begin. Notice the crucial role of basic research, using experimental animals, in this process. Virtually every prescription drug on the market owes its existence to such research.

🖌 🛛 CHECKPOINTS

- 1. How has the study of physiology aided, and been aided by, the study of diseases?
- **2a.** Describe the steps involved in the scientific method. What would qualify a statement as unscientific?
- **2b.** Describe the different types of trials a new drug must undergo before it is "ready for market."

1.2 HOMEOSTASIS AND FEEDBACK CONTROL

The regulatory mechanisms of the body can be understood in terms of a single shared function: that of maintaining constancy of the internal environment. A state of relative constancy of the internal environment is known as homeostasis, maintained by negative feedback loops.

LEARNING OUTCOMES

After studying this section, you should be able to:

- Define homeostasis, and identify the components of negative feedback loops.
- **4.** Explain the role of antagonistic effectors in maintaining homeostasis, and the nature of positive feedback loops.
- Give examples of how negative feedback loops involving the nervous and endocrine systems help to maintain homeostasis.

History of Physiology

The Greek philosopher Aristotle (384–322 B.C.) speculated on the function of the human body, but another ancient Greek, Erasistratus (304–250? B.C.), is considered to be the first to study physiology because he attempted to apply physical laws to understand human function. Galen (A.D. 130–201) wrote widely on the subject and was considered the supreme authority until the Renaissance. Physiology became a fully experimental science with the revolutionary work of the English physician William Harvey (1578–1657), who demonstrated that the heart pumps blood through a closed system of vessels.

However, the originator of modern physiology is the French physiologist Claude Bernard (1813–1878), who observed that the *milieu intérieur* (internal environment) remains remarkably constant despite changing conditions in the external environment. In a book entitled *The Wisdom of the Body*, published in 1932, the American physiologist Walter Cannon (1871–1945) coined the term **homeostasis** to describe this internal constancy. Cannon further suggested that the many mechanisms of physiological regulation have but one purpose—the maintenance of internal constancy.

Most of our present knowledge of human physiology has been gained in the twentieth century. However, new knowledge in the twenty-first century is being added at an ever more rapid pace, fueled in more recent decades by the revolutionary growth of molecular genetics and its associated biotechnologies, and by the availability of more powerful computers and other equipment. A very brief history of twentieth- and twentyfirst-century physiology, limited by space to only two citations per decade, is provided in table 1.1.

Most of the citations in table 1.1 indicate the winners of Nobel prizes. The **Nobel Prize in Physiology or Medicine** (a single prize category) was first awarded in 1901 to Emil Adolf von Behring, a pioneer in immunology who coined the term

Table 1.1 | History of Twentieth- and Twenty-First-Century Physiology(two citations per decade)

1900	Karl Landsteiner discovers the A, B, and O blood groups.
1904	Ivan Pavlov wins the Nobel Prize for his work on the physiology of digestion.
1910	Sir Henry Dale describes properties of histamine.
1918	Earnest Starling describes how the force of the heart's contraction relates to the amount of blood in it.
1921	John Langley describes the functions of the autonomic nervous system.
1923	Sir Frederick Banting, Charles Best, and John Macleod win the Nobel Prize for the discovery of insulin.
1932	Sir Charles Sherrington and Lord Edgar Adrian win the Nobel Prize for discoveries related to the functions of neurons.
1936	Sir Henry Dale and Otto Loewi win the Nobel Prize for the discovery of acetylcholine in synaptic transmission.
1939–47	Albert von Szent-Györgyi explains the role of ATP and contributes to the understanding of actin and myosin in muscle contraction.
1949	Hans Selye discovers the common physiological responses to stress.
1953	Sir Hans Krebs wins the Nobel Prize for his discovery of the citric acid cycle.
1954	Hugh Huxley, Jean Hanson, R. Niedergerde, and Andrew Huxley propose the sliding filament theory of muscle contraction.
1962	Francis Crick, James Watson, and Maurice Wilkins win the Nobel Prize for determining the structure of DNA.
1963	Sir John Eccles, Sir Alan Hodgkin, and Sir Andrew Huxley win the Nobel Prize for their discoveries relating to the nerve impulse.
1971	Earl Sutherland wins the Nobel Prize for his discovery of the mechanism of hormone action.
1977	Roger Guillemin and Andrew Schally win the Nobel Prize for discoveries of the brain's production of peptide hormone.
1981	Roger Sperry wins the Nobel Prize for his discoveries regarding the specializations of the right and left cerebral hemispheres.
1986	Stanley Cohen and Rita Levi-Montalcini win the Nobel Prize for their discoveries of growth factors regulating the nervous system.
1994	Alfred Gilman and Martin Rodbell win the Nobel Prize for their discovery of the functions of G-proteins in signal transduction in cells.
1998	Robert Furchgott, Louis Ignarro, and Ferid Murad win the Nobel Prize for discovering the role of nitric oxide as a signaling molecule in the cardiovascular system.
2004	Linda B. Buck and Richard Axel win the Nobel Prize for their discoveries of odorant receptors and the organization of the olfactory system.
2006	Andrew Z. Fine and Craig C. Mello win the Noble Prize for their discovery of RNA interference by short, double-stranded RNA molecules.

antibody and whose many other discoveries included the use of serum (containing antibodies) to treat diphtheria. Many scientists who might deserve a Nobel Prize never receive one, and the prizes are given for particular achievements and not others (Einstein didn't win his Nobel Prize in Physics for relativity, for example) and are often awarded many years after the discoveries were made. Nevertheless, the awarding of the Nobel Prize in Physiology or Medicine each year is a celebrated event in the biomedical community, and the awards can be a useful yardstick for tracking the course of physiological research over time.

Negative Feedback Loops

The concept of homeostasis has been of immense value in the study of physiology because it allows diverse regulatory mechanisms to be understood in terms of their "why" as well as their "how." The concept of homeostasis also provides a major foundation for medical diagnostic procedures. When a particular measurement of the internal environment, such as a blood measurement (table 1.2), deviates significantly from the normal range of values, it can be concluded that homeostasis is not being maintained and that the person is sick. A number of such measurements, combined with clinical observations, may allow the particular defective mechanism to be identified.

In order for internal constancy to be maintained, changes in the body must stimulate **sensors** that can send information to an **integrating center**. This allows the integrating center to detect changes from a **set point**. The set point is analogous to the temperature set on a house thermostat. In a similar manner, there is a set point for body temperature, blood glucose concentration, the tension on a tendon, and so on. The integrating center is often a particular region of the brain or spinal cord, but it can also be a group of cells in an endocrine gland. A number of

Table 1.2 | Approximate Normal Rangesfor Measurements of Some FastingBlood Values

Measurement	Normal Range	
Arterial pH	7.35–7.45	
Bicarbonate	24–28 mEq/L	
Sodium	135–145 mEq/L	
Calcium	4.5–5.5 mEq/L	
Oxygen content	17.2–22.0 ml/100 ml	
Urea	12-35 mg/100 ml	
Amino acids	3.3–5.1 mg/100 ml	
Protein	6.5–8.0 g/100 ml	
Total lipids	400–800 mg/100 ml	
Glucose	75–110 mg/100 ml	

different sensors may send information to a particular integrating center, which can then integrate this information and direct the responses of **effectors**—generally muscles or glands. The integrating center may cause increases or decreases in effector action to counter the deviations from the set point and defend homeostasis.

The thermostat of a house can serve as a simple example. Suppose you set the thermostat at a set point of 70° F. If the temperature in the house rises sufficiently above the set point, a sensor connected to an integrating center within the thermostat will detect that deviation and turn on the air conditioner (the effector in this example). The air conditioner will turn off when the room temperature falls and the thermostat no longer detects a deviation from the set-point temperature. However, this simple example gives a wrong impression: the effectors in the body are generally increased or decreased in activity, *not* just turned on or off. Because of this, negative feedback control in the body works far more efficiently than does a house thermostat.

If the body temperature exceeds the set point of 37° C, sensors in a part of the brain detect this deviation and, acting via an integrating center (also in the brain), stimulate activities of effectors (including sweat glands) that lower the temperature. For another example, if the blood glucose concentration falls below normal, the effectors act to increase the blood glucose. One can think of the effectors as "defending" the set points against deviations. Because the activity of the effectors is influenced by the effects they produce, and because this regulation is in a negative, or reverse, direction, this type of control system is known as a **negative feedback loop** (fig. 1.1). (Notice that in figure 1.1 and in all subsequent figures, negative feedback is indicated by a dashed line and a negative sign.)






Figure 1.1 A rise in some factor of the internal environment (\uparrow X) is detected by a sensor. This information is relayed to an integrating center, which causes an effector to produce a change (1) in the opposite direction (\downarrow X). The initial deviation is thus reversed (2), completing a negative feedback loop (shown by the dashed arrow and negative sign). The numbers indicate the sequence of changes.





The nature of the negative feedback loop can be understood by again referring to the analogy of the thermostat and air conditioner. After the air conditioner has been on for some time, the room temperature may fall significantly below the set point of the thermostat. When this occurs, the air conditioner will be turned off. The effector (air conditioner) is turned on by a high temperature and, when activated, produces a negative change (lowering of the temperature) that ultimately causes the effector to be turned off. In this way, constancy is maintained.

It is important to realize that these negative feedback loops are continuous, ongoing processes. Thus, a particular nerve fiber that is part of an effector mechanism may always display some activity, and a particular hormone that is part of another effector mechanism may always be present in the blood. The nerve activity and hormone concentration may decrease in response to deviations of the internal environment in one direction (fig. 1.1), or they may increase in response to deviations in the opposite direction (fig. 1.2). Changes from the normal range in either direction are thus compensated for by reverse changes in effector activity.

Because negative feedback loops respond after deviations from the set point have stimulated sensors, the internal environment is never absolutely constant. Homeostasis is best conceived as a state of **dynamic constancy** in which conditions are stabilized above and below the set point. These conditions can be measured quantitatively, in degrees Celsius for body temperature, for example, or in milligrams per deciliter (one-tenth of a liter) for blood glucose. The set point can be taken as the average value within the normal range of measurements (fig. 1.3).

Antagonistic Effectors

Most factors in the internal environment are controlled by several effectors, which often have antagonistic actions. Control by antagonistic effectors is sometimes described as "push-pull," where the increasing activity of one effector is




Figure 1.3 Negative feedback loops maintain a state of dynamic constancy within the internal environment. The completion of the negative feedback loop is indicated by negative signs.




Figure 1.4 How body temperature is maintained within the normal range. The body temperature normally has a set point of 37° C. This is maintained, in part, by two antagonistic mechanisms—shivering and sweating. Shivering is induced when the body temperature falls too low, and it gradually subsides as the temperature rises. Sweating occurs when the body temperature is too high, and it diminishes as the temperature falls. Most aspects of the internal environment are regulated by the antagonistic actions of different effector mechanisms.

See the Test Your Quantitative Ability section of the Review Activities at the end of this chapter.

accompanied by decreasing activity of an antagonistic effector. This affords a finer degree of control than could be achieved by simply switching one effector on and off.

Room temperature can be maintained, for example, by simply turning an air conditioner on and off, or by just turning a heater on and off. A much more stable temperature, however, can be achieved if the air conditioner and heater are both controlled by a thermostat. Then the heater is turned on when the air conditioner is turned off, and vice versa. Normal body temperature is maintained about a set point of 37° C by the antagonistic effects of sweating, shivering, and other mechanisms (fig. 1.4).

The blood concentrations of glucose, calcium, and other substances are regulated by negative feedback loops involving hormones that promote opposite effects. Insulin, for example, lowers blood glucose, and other hormones raise the blood glucose concentration. The heart rate, similarly, is controlled by nerve fibers that produce opposite effects: stimulation of one group of nerve fibers increases heart rate; stimulation of another group slows the heart rate.

Quantitative Measurements

In order to study physiological mechanisms, scientists must measure specific values and mathematically determine such statistics as their normal range, their averages, and their




Figure 1.5 Homeostasis of the blood glucose

concentration. Average blood glucose concentrations of five healthy individuals are graphed before and after a rapid intravenous injection of insulin. The "0" indicates the time of the injection. The blood glucose concentration is first lowered by the insulin injection, but is then raised back to the normal range (by hormones antagonistic to insulin that stimulate the liver to secrete glucose into the blood). Homeostasis of blood glucose is maintained by the antagonistic actions of insulin and several other hormones.

deviations from the average (which can represent the set point). For these and other reasons, quantitative measurements are basic to the science of physiology. One example of this, and of the actions of antagonistic mechanisms in maintaining homeostasis, is shown in figure 1.5. Blood glucose concentrations were measured in five healthy people before and after an injection of insulin, a hormone that acts to lower the blood glucose concentration. A graph of the data reveals that the blood glucose concentration decreased rapidly but was brought back up to normal levels within 80 minutes after the injection. This demonstrates that negative feedback mechanisms acted to restore homeostasis in this experiment. These mechanisms involve the action of hormones whose effects are antagonistic to that of insulinthat is, they promote the secretion of glucose from the liver (see chapter 19).

Positive Feedback

Constancy of the internal environment is maintained by effectors that act to compensate for the change that served as the stimulus for their activation; in short, by negative feedback loops. A thermostat, for example, maintains a constant temperature by increasing heat production when it is cold and decreasing heat production when it is warm. The opposite occurs during **positive feedback**—in this case, the action of effectors *amplifies* those changes that stimulated the effectors. A thermostat that works by positive feedback, for example, would increase heat production in response to a rise in temperature.

It is clear that homeostasis must ultimately be maintained by negative rather than by positive feedback mechanisms. The effectiveness of some negative feedback loops, however, is increased by positive feedback mechanisms that amplify the actions of a negative feedback response. Blood clotting, for example, occurs as a result of a sequential activation of clotting factors; the activation of one clotting factor results in activation of many in a positive feedback cascade. In this way, a single change is amplified to produce a blood clot. Formation of the clot, however, can prevent further loss of blood, and thus represents the completion of a negative feedback loop that restores homeostasis.

Two other examples of positive feedback in the body are both related to the female reproductive system. One of these examples occurs when estrogen, secreted by the ovaries, stimulates the women's pituitary gland to secrete LH (luteinizing hormone). This stimulatory, positive feedback effect creates an "LH surge" (very rapid rise in blood LH concentrations) that triggers ovulation. Interestingly, estrogen secretion after ovulation has an inhibitory, negative feedback, effect on LH secretion (this is the physiological basis for the birth control pill, discussed in chapter 20). Another example of positive feedback is contraction of the uterus during childbirth (parturition). Contraction of the uterus is stimulated by the pituitary hormone oxytocin, and the secretion of oxytocin is increased by sensory feedback from contractions of the uterus during labor. The strength of uterine contractions during labor is thus increased through positive feedback. The mechanisms involved in labor are discussed in more detail in chapter 20 (see fig. 20.50).

Neural and Endocrine Regulation

Homeostasis is maintained by two general categories of regulatory mechanisms: (1) those that are **intrinsic**, or "built into" the organs being regulated (such as molecules produced in the walls of blood vessels that cause vessel dilation or constriction); and (2) those that are **extrinsic**, as in regulation of an organ by the nervous and endocrine systems. The endocrine system functions closely with the nervous system in regulating and integrating body processes and maintaining homeostasis. The nervous system controls the secretion of many endocrine glands, and some hormones in turn affect the function of the nervous system. Together, the nervous and endocrine systems of the body.

Regulation by the endocrine system is achieved by the secretion of chemical regulators called **hormones** into the blood, which carries the hormones to all organs in the body. Only specific organs can respond to a particular hormone, however; these are known as the **target organs** of that hormone.

Nerve fibers are said to *innervate* the organs that they regulate. When stimulated, these fibers produce electrochemical nerve impulses that are conducted from the origin of the fiber to its terminals in the target organ innervated by the fiber. These target organs can be muscles or glands that may function as effectors in the maintenance of homeostasis.




Figure 1.6 Negative feedback control of blood pressure. Blood pressure influences the activity of sensory neurons from the blood pressure receptors (sensors); a rise in pressure increases the firing rate, and a fall in pressure decreases the firing rate of nerve impulses. When a person stands up from a lying-down position, the blood pressure momentarily falls. The resulting decreased firing rate of nerve impulses in sensory neurons affects the medulla oblongata of the brain (the integrating center). This causes the motor nerves to the heart (effector) to increase the heart rate, helping to raise the blood pressure.

For example, we have negative feedback loops that help maintain homeostasis of arterial blood pressure, in part by adjusting the heart rate. If everything else is equal, blood pressure is lowered by a decreased heart rate and raised by an increased heart rate. This is accomplished by regulating the activity of the autonomic nervous system, as will be discussed in later chapters. Thus, a fall in blood pressure—produced daily as we go from a lying to a standing position—is compensated by a faster heart rate (fig. 1.6). As a consequence of this negative feedback loop, our heart rate varies as we go through our day, speeding up and slowing down, so that we can maintain homeostasis of blood pressure and keep it within normal limits.

Feedback Control of Hormone Secretion

The nature of the endocrine glands, the interaction of the nervous and endocrine systems, and the actions of hormones will be discussed in detail in later chapters. For now, it is sufficient to describe the regulation of hormone secretion very broadly, because it so superbly illustrates the principles of homeostasis and negative feedback regulation.

Hormones are secreted in response to specific chemical stimuli. A rise in the plasma glucose concentration, for example, stimulates insulin secretion from structures in the pancreas known as the *pancreatic islets*, or *islets of Langerhans*. Hormones are also secreted in response to nerve stimulation and stimulation by other hormones.

The secretion of a hormone can be inhibited by its own effects in a negative feedback manner. Insulin, as previously described, produces a lowering of blood glucose. Because a rise in blood glucose stimulates insulin secretion, a lowering of blood glucose caused by insulin's action inhibits further insulin secretion. This closed-loop control system is called **negative feedback inhibition** (fig. 1.7a).

Homeostasis of blood glucose is too important—the brain uses blood glucose as its primary source of energy—to entrust to the regulation of only one hormone, insulin. So, when blood glucose falls during fasting, several mechanisms prevent it from falling too far (fig. 1.7b). First, insulin secretion decreases, preventing muscle, liver, and adipose cells from taking too much glucose from the blood. Second, the secretion of a hormone antagonistic to insulin, called *glucagon*, increases. Glucagon stimulates processes in the liver (breakdown of a stored, starchlike molecule called glycogen; chapter 2, section 2.2) that cause it to secrete glucose into the blood. Through these and other antagonistic negative feedback mechanisms, the blood glucose is maintained within a homeostatic range.




Figure 1.7 Negative feedback control of blood glucose. (a) The rise in blood glucose that occurs after eating carbohydrates is corrected by the action of insulin, which is secreted in increasing amounts at that time. (b) During fasting, when blood glucose falls, insulin secretion is inhibited and the secretion of an antagonistic hormone, glucagon, is increased. This stimulates the liver to secrete glucose into the blood, helping to prevent blood glucose from continuing to fall. In this way, blood glucose concentrations are maintained within a homeostatic range following eating and during fasting.

Clinical Investigation

CLUES

Clinical Investigation Clues are placed immediately following the text information that pertains to the Clinical Investigation for the chapter. Use these to solve the medical mystery—if you need to, re-read the information preceding the "Clues." You can check your answers against the Clinical Investigation Summaries at the end of the chapters. In this case, Linda had a normal resting body temperature and a normal fasting glucose concentration, suggesting that homeostasis of these values was being maintained.

C H E C K P O I N T S

- **3a.** Define *homeostasis* and describe how this concept can be used to explain physiological control mechanisms.
- **3b.** Define *negative feedback* and explain how it contributes to homeostasis. Illustrate this concept by drawing and labeling a negative feedback loop.
- 4. Describe *positive feedback* and explain how this process functions in the body.
- **5.** Explain how the secretion of a hormone is controlled by negative feedback inhibition. Use the control of insulin secretion as an example.

1.3 THE PRIMARY TISSUES

The organs of the body are composed of four different primary tissues, each of which has its own characteristic structure and function. The activities and interactions of these tissues determine the physiology of the organs.

LEARNING OUTCOMES

After studying this section, you should be able to:

- 6. Distinguish the primary tissues and their subtypes.
- **7.** Relate the structure of the primary tissues to their functions.

Although physiology is the study of function, it is difficult to properly understand the function of the body without some knowledge of its anatomy, particularly at a microscopic level. Microscopic anatomy constitutes a field of study known as *histology*. The anatomy and histology of specific organs will be discussed together with their functions in later chapters. In this section, the common "fabric" of all organs is described.

Cells are the basic units of structure and function in the body. Cells that have similar functions are grouped into categories called *tissues*. The entire body is composed of only four major types of tissues. These **primary tissues** are (1) muscle,

(2) nervous, (3) epithelial, and (4) connective tissues. Groupings of these four primary tissues into anatomical and functional units are called **organs**. Organs, in turn, may be grouped together by common functions into **systems**. The systems of the body act in a coordinated fashion to maintain the entire organism.

Muscle Tissue

Muscle tissue is specialized for contraction. There are three types of muscle tissue: **skeletal, cardiac,** and **smooth.** Skeletal muscle is often called *voluntary muscle* because its contraction is consciously controlled. Both skeletal and cardiac muscles are **striated;** they have striations, or stripes, that extend across the width of the muscle cell (figs. 1.8 and 1.9). These striations are produced by a characteristic arrangement of contractile proteins, and for this reason skeletal and cardiac muscle have similar mechanisms of contraction. Smooth muscle (fig. 1.10) lacks these striations and has a different mechanism of contraction.

Skeletal Muscle

Skeletal muscles are generally attached to bones at both ends by means of tendons; hence, contraction produces movements of the skeleton. There are exceptions to this pattern, however. The tongue, superior portion of the esophagus, anal sphincter, and diaphragm are also composed of skeletal muscle, but they do not cause movements of the skeleton.

Beginning at about the fourth week of embryonic development, separate cells called *myoblasts* fuse together to form **skeletal muscle fibers**, or **myofibers** (from the Greek myos = muscle). Although myofibers are often referred to as skeletal muscle cells, each is actually a *syncytium*, or multinucleate mass formed from the union of separate cells. Despite their unique origin and structure, each myofiber contains mitochondria and other organelles (described in chapter 3) common to all cells.

The muscle fibers within a skeletal muscle are arranged in bundles, and within these bundles the fibers extend in parallel from one end of the bundle to the other. The parallel arrangement of muscle fibers (fig. 1.8) allows each fiber to be controlled individually: one can thus contract fewer or more muscle fibers and, in this way, vary the strength of contraction of the whole muscle. The ability to vary, or "grade," the strength of skeletal muscle contraction is needed for precise control of skeletal movements.

Cardiac Muscle

Although cardiac muscle is striated, it differs markedly from skeletal muscle in appearance. Cardiac muscle is found only in the heart where the **myocardial cells** are short, branched, and intimately interconnected to form a continuous fabric. Special areas of contact between adjacent cells stain darkly to show




Figure 1.8 Skeletal muscle fibers showing the characteristic light and dark cross striations. Because of this feature, skeletal muscle is also called striated muscle. AP R




Figure 1.9 Human cardiac muscle. Notice the striated appearance and dark-staining intercalated discs. AP**|R**




Figure 1.10 A photomicrograph of smooth muscle cells. Notice that these cells contain single, centrally located nuclei and lack striations. AP **R**

intercalated discs (fig. 1.9), which are characteristic of heart muscle.

The intercalated discs couple myocardial cells together mechanically and electrically. Unlike skeletal muscles, therefore, the heart cannot produce a graded contraction by varying the number of cells stimulated to contract. Because of the way the heart is constructed, the stimulation of one myocardial cell results in the stimulation of all other cells in the mass and a "wholehearted" contraction.

Smooth Muscle

As implied by the name, smooth muscle cells (fig. 1.10) do not have the striations characteristic of skeletal and cardiac muscle. Smooth muscle is found in the digestive tract, blood vessels, bronchioles (small air passages in the lungs), and the ducts of the urinary and reproductive systems. Circular arrangements of smooth muscle in these organs produce constriction of the *lumen* (cavity) when the muscle cells contract. The digestive tract also contains longitudinally arranged layers of smooth muscle. *Peristalsis* is the coordinated wavelike contractions of the circular and longitudinal smooth muscle layers that push food from the oral to the anal end of the digestive tract.

The three types of muscle tissue are discussed further in chapter 12.

Nervous Tissue

Nervous tissue consists of nerve cells, or **neurons**, which are specialized for the generation and conduction of electrical events, and **neuroglial** (or **glial**) **cells.** Neuroglial cells provide the neurons with structural support and perform a variety of functions that are needed for the normal physiology of the nervous system.

Each neuron consists of three parts: (1) a *cell body*, (2) *dendrites*, and (3) an *axon* (fig. 1.11). The cell body contains the nucleus and serves as the metabolic center of the cell. The dendrites (literally, "branches") are highly branched cytoplasmic extensions of the cell body that receive input from other neurons or from receptor cells. The axon is a single cytoplasmic extension of the cell body that can be quite long (up to a few feet in length). It is specialized for conducting nerve




Figure 1.11 A photomicrograph of nerve tissue. A single neuron and numerous smaller supporting cells can be seen. APIR

impulses from the cell body to another neuron or to an effector (muscle or gland) cell.

The neuroglial cells do not conduct impulses but instead serve to bind neurons together, modify the extracellular environment of the nervous system, and influence the nourishment and electrical activity of neurons. In recent years, neuroglial cells have been shown to cooperate with neurons in chemical neurotransmission (chapter 7), and to have many other roles in the normal physiology (as well as disease processes) of the brain and spinal cord. Neuroglial cells are about five times more abundant than neurons in the nervous system and, unlike neurons, maintain a limited ability to divide by mitosis throughout life.

Neurons and neuroglial cells are discussed in detail in chapter 7.

Epithelial Tissue

Epithelial tissue consists of cells that form **membranes**, which cover and line the body surfaces, and of **glands**, which are derived from these membranes. There are two categories of glands. *Exocrine glands* (from the Greek *exo* = outside) secrete chemicals through a duct that leads to the outside of a membrane, and thus to the outside of a body surface. *Endocrine glands* (from the Greek *endon* = within) secrete chemicals called *hormones* into the blood. Endocrine glands are discussed in chapter 11.

Epithelial Membranes


Epithelial membranes are classified according to the number of their layers and the shape of the cells in the upper layer (table 1.3). Epithelial cells that are flattened in shape are **squamous;** those that are as wide as they are tall are **cuboidal;** and those that are taller than they are wide are **columnar** (fig. 1.12a-c). Those epithelial membranes that are only one cell layer thick are known as **simple membranes;** those that are composed of a number of layers are **stratified membranes.**


Epithelial membranes cover all body surfaces and line the cavity (lumen) of every hollow organ. Thus, epithelial membranes provide a barrier between the external environment and the internal environment of the body. Stratified epithelial membranes are specialized to provide protection. Simple epithelial membranes, in contrast, provide little protection; instead, they are specialized for transport of substances between the internal and external environments. In order for a substance to get into the body, it must pass through an epithelial membrane, and simple epithelia are specialized for this function. For example, a simple squamous epithelium in the lungs allows the rapid passage of oxygen and carbon dioxide between the air (external environment) and blood (internal environment). A simple columnar epithelium in the small intestine, as another example, allows digestion products to pass from the intestinal lumen (external environment) to the blood (internal environment).

Dispersed among the columnar epithelial cells are specialized unicellular glands called *goblet cells* that secrete mucus. The columnar epithelial cells in the uterine (fallopian) tubes of


Table 1.3 | Summary of Epithelial Membranes

Туре	Structure and Function	Location
Simple Epithelia	Single layer of cells; function varies with type	Covering visceral organs; linings of body cavities, tubes, and ducts
Simple squamous epithelium	Single layer of flattened, tightly bound cells; diffusion and filtration	Capillary walls; pulmonary alveoli of lungs; covering visceral organs; linings of body cavities
Simple cuboidal epithelium	Single layer of cube-shaped cells; excretion, secretion, or absorption	Surface of ovaries; linings of kidney tubules, salivary ducts, and pancreatic ducts
Simple columnar epithelium	Single layer of nonciliated, tall, column-shaped cells; protection, secretion, and absorption	Lining of most of digestive tract
Simple ciliated columnar epithelium	Single layer of ciliated, column-shaped cells; transportive role through ciliary motion	Lining of uterine tubes
Pseudostratified ciliated columnar epithelium	Single layer of ciliated, irregularly shaped cells; many goblet cells; protection, secretion, ciliary movement	Lining of respiratory passageways
Stratified Epithelia	Two or more layers of cells; function varies with type	Epidermal layer of skin; linings of body openings, ducts, and urinary bladder
Stratified squamous epithelium (keratinized)	Numerous layers containing keratin, with outer layers flattened and dead; protection	Epidermis of skin
Stratified squamous epithelium (nonkeratinized)	Numerous layers lacking keratin, with outer layers moistened and alive; protection and pliability	Linings of oral and nasal cavities, vagina, and anal canal
Stratified cuboidal epithelium	Usually two layers of cube-shaped cells; strengthening of luminal walls	Large ducts of sweat glands, salivary glands, and pancreas
Transitional epithelium	Numerous layers of rounded, nonkeratinized cells; distension	Walls of ureters, part of urethra, and urinary bladder





(c)






Figure 1.13 A stratified squamous nonkeratinized epithelial membrane. This is a photomicrograph (*a*) and illustration (*b*) of the epithelial lining of the vagina. AP|R

females and in the respiratory passages contain numerous *cilia* (hairlike structures, described in chapter 3) that can move in a coordinated fashion and aid the functions of these organs.

The epithelial lining of the esophagus and vagina that provides protection for these organs is a stratified squamous epithelium (fig. 1.13). This is a *nonkeratinized* membrane, and all layers consist of living cells. The *epidermis* of the skin, by contrast, is *keratinized*, or *cornified* (fig. 1.14). Because the epidermis is dry and exposed to the potentially desiccating effects of the air, the surface is covered with dead cells that are filled with a water-resistant protein known as *keratin*. This protective layer is constantly flaked off from the surface of the skin and therefore must be constantly replaced by the division of cells in the deeper layers of the epidermis.

The constant loss and renewal of cells is characteristic of epithelial membranes. The entire epidermis is completely replaced every two weeks; the stomach lining is renewed every two to three days. Examination of the cells that are lost, or "exfoliated," from the outer layer of epithelium lining the female reproductive tract is a common procedure in gynecology (as in the Pap smear).

In order to form a strong membrane that is effective as a barrier at the body surfaces, epithelial cells are very closely packed and are joined together by structures collectively called **junctional complexes** (chapter 6; see fig. 6.22). There is no room for blood vessels between adjacent epithelial cells. The epithelium must therefore receive nourishment from the tissue beneath, which has large intercellular spaces that can accommodate blood vessels and nerves. This underlying tissue is called *connective tissue*. Epithelial membranes are attached to the underlying connective tissue by a layer of proteins and polysaccharides known as the **basement membrane**. This layer can be observed only under the microscope using specialized staining techniques.

Basement membranes are believed to induce a polarity to the cells of epithelial membranes; that is, the top (apical) portion of epithelial cells has different structural and functional components than the bottom (basal) portion. This is important in many physiological processes. For example, substances are transported in specific directions across simple epithelial membranes (discussed in chapter 6; see fig. 6.21). In stratified membranes, only the basal (bottom) layer of cells is on the basement membrane, and it is these cells that undergo mitosis to form new epithelial




Figure 1.14 The epidermis is a stratified, squamous, keratinized epithelium. The upper cell layers are dead and impregnated with the protein keratin, producing a cornified epithelial membrane, which is supported by layers of living cells. The epidermis is nourished by blood vessels located in the loose connective tissue of the dermis. **AP**

cells to replace those lost from the top. Scientists recently demonstrated that when these basal cells divide, one of the daughter cells is attached to the basement membrane (renewing the basal cell population), while the other is not. The daughter cell that is "unstuck" from the basement membrane differentiates and migrates upward in the stratified epithelium.

CLINICAL APPLICATION

Basement membranes consist primarily of the structural protein known as *collagen* (see fig. 1.17). The type of collagen in basement membranes is a large protein assembled from six different subunits. **Alport syndrome** is a genetic disorder of the collagen subunits that, among other problems, results in damage to the glomeruli (the filtering units) of the kidneys. This is one of the most common causes of kidney failure. In **Goodpasture disease**, the collagen in the basement membranes of the glomeruli and the lungs is attacked by the person's own antibodies, leading to both kidney and lung disease.

Exfoliative cytology is the collection and examination of epithelial cells that are shed and collected by mechanical scraping of the membranes, washing of the membranes, or aspiration of body fluids containing the shed cells. Microscopic examination of these *desquamated* (shed) cells, for example in a **Pap smear,** may reveal a malignancy.

Exocrine Glands

Exocrine glands are derived from cells of epithelial membranes. The secretions of these cells are passed to the outside of the epithelial membranes (and hence to the surface of the body) through *ducts*. This is in contrast to *endocrine glands*, which lack ducts and which therefore secrete into capillaries within the body (fig. 1.15). The structure of endocrine glands will be described in chapter 11.

The secretory units of exocrine glands may be simple tubes, or they may be modified to form clusters of units around branched ducts (fig. 1.16). These clusters, or **acini**, are often surrounded by tentaclelike extensions of *myoepithelial cells* that contract and squeeze the secretions through the ducts. The rate of secretion and the action of myoepithelial cells are subject to neural and endocrine regulation.

Examples of exocrine glands in the skin include the lacrimal (tear) glands, sebaceous glands (which secrete oily sebum into hair follicles), and sweat glands. There are two types of sweat glands. The more numerous, the *eccrine* (or *merocrine*) *sweat glands*, secrete a dilute salt solution that serves in thermoregulation (evaporation cools the skin). The *apocrine sweat glands*, located in the axillae (underarms) and pubic region, secrete a protein-rich fluid. This provides nourishment for bacteria that produce the characteristic odor of this type of sweat.

All of the glands that secrete into the digestive tract are also exocrine. This is because the lumen of the digestive tract is a part of the external environment, and secretions of these glands go to the outside of the membrane that lines this tract. Mucous glands are located throughout the length of the digestive tract. Other relatively simple glands of the tract include salivary glands, gastric glands, and simple tubular glands in the intestine.

The *liver* and *pancreas* are exocrine (as well as endocrine) glands, derived embryologically from the digestive tract. The exocrine secretion of the pancreas—pancreatic juice—contains digestive enzymes and bicarbonate and is secreted into the small intestine via the pancreatic duct. The liver produces and secretes bile (an emulsifier of fat) into the small intestine via the gallbladder and bile duct.




Figure 1.15 The formation of exocrine and endocrine glands from epithelial membranes. Note that exocrine glands retain a duct that can carry their secretion to the surface of the epithelial membrane, whereas endocrine glands are ductless.




Figure 1.16 The structure of exocrine glands. Exocrine glands may be simple invaginations of epithelial membranes, or they may be more complex derivatives. APR

Exocrine glands are also prominent in the reproductive system. The female reproductive tract contains numerous mucussecreting exocrine glands. The male accessory sex organs—the *prostate* and *seminal vesicles*—are exocrine glands that contribute to semen. The testes and ovaries (the gonads) are both endocrine and exocrine glands. They are endocrine because they secrete sex steroid hormones into the blood; they are exocrine because they release gametes (ova and sperm) into the reproductive tracts.

Connective Tissue

Connective tissue is characterized by large amounts of extracellular material between the different types of connective tissue cells. The extracellular material, called the connective tissue *matrix*, varies in the four primary types of connective tissues: (1) connective tissue proper; (2) cartilage; (3) bone; and (4) blood. **Blood** is classified as a type of connective tissue because about half its volume is an extracellular fluid, the blood plasma (chapter 13, section 13.1).

Connective tissue proper, in which the matrix consists of protein fibers and a proteinaceous, gel-like ground substance, is divided into subtypes. In loose connective tissue (also called areolar connective tissue), protein fibers composed of collagen (collagenous fibers) are scattered loosely in the ground substance (fig. 1.17), which provides space for the presence of blood vessels, nerve fibers, and other structures (see the dermis of the skin, shown in fig. 1.14, as an example). Dense regular connective tissues are those in which collagenous fibers are oriented parallel to each other and densely packed in the extracellular matrix, leaving little room for cells and ground substance (fig. 1.18). Examples of dense regular connective tissues include tendons (connecting bone to bone) and ligaments (connecting bones together at joints). Dense irregular connective tissues, forming tough capsules and sheaths around organs, contain densely packed collagenous fibers arranged in various orientations that resist forces applied from different directions.

Adipose tissue is a specialized type of loose connective tissue. In each adipose cell, or *adipocyte*, the cytoplasm is stretched around a central globule of fat (fig. 1.19). The synthesis and breakdown of fat are accomplished by enzymes within the cytoplasm of the adipocytes.

Cartilage consists of cells, called *chondrocytes*, surrounded by a semisolid ground substance that imparts elastic properties




Figure 1.17 Loose connective tissue. This illustration shows the cells and protein fibers characteristic of connective tissue proper. The ground substance is the extracellular background material, against which the different protein fibers can be seen. The macrophage is a phagocytic connective tissue cell, which can be derived from monocytes (a type of white blood cell).

to the tissue. Cartilage is a type of supportive and protective tissue commonly called "gristle." It forms the precursor to many bones that develop in the fetus and persists at the articular (joint) surfaces on the bones at all movable joints in adults.

Bone is produced as concentric layers, or *lamellae*, of calcified material laid around blood vessels. The bone-forming cells, or *osteoblasts*, surrounded by their calcified products, become trapped within cavities called *lacunae*. The trapped cells, which are now called *osteocytes*, remain




Figure 1.18 Dense regular connective tissue. In this photomicrograph, the collagen fibers in a tendon are packaged densely into parallel groups. The ground substance is in the tiny spaces between the collagen fibers. AP




Figure 1.19 Adipose tissue. Each adipocyte contains a large, central globule of fat surrounded by the cytoplasm of the adipocyte. (*a*) Photomicrograph and (*b*) illustration of adipose tissue. AP **R**

alive because they are nourished by "lifelines" of cytoplasm that extend from the cells to the blood vessels in *canaliculi* (little canals). The blood vessels lie within central canals, surrounded by concentric rings of bone lamellae with their trapped osteocytes. These units of bone structure are called *osteons*, or *haversian systems* (fig. 1.20).




Figure 1.20 The structure of bone. (*a*) A diagram of a long bone, (*b*) a photomicrograph showing osteons (haversian systems), and (*c*) a diagram of osteons. Within each central canal, an artery (red), a vein (blue), and a nerve (yellow) is illustrated. **APR**




Figure 1.21 A cross section of a tooth showing pulp, dentin, and enamel. The root of the tooth is covered by cementum, a calcified connective tissue that helps to anchor the tooth in its bony socket.

The *dentin* of a tooth (fig. 1.21) is similar in composition to bone, but the cells that form this calcified tissue are located in the pulp (composed of loose connective tissue). These cells send cytoplasmic extensions, called *dentinal tubules*, into the dentin. Dentin, like bone, is thus a living tissue that can be remodeled in response to stresses. The cells that form the outer *enamel* of a tooth, by contrast, are lost as the tooth erupts. Enamel is a highly calcified material, harder than bone or dentin, that cannot be regenerated; artificial "fillings" are therefore required to patch holes in the enamel.

CHECKPOINTS

- **6a.** List the four primary tissues and describe the distinguishing features of each type.
- 6b. Compare and contrast the three types of muscle tissue.
- **6c.** Describe the different types of epithelial membranes
- and state their locations in the body. **7a.** Explain why exocrine and endocrine glands are
- considered epithelial tissues and distinguish between these two types of glands.
- **7b.** Describe the different types of connective tissues and explain how they differ from one another in their content of extracellular material.

1.4 ORGANS AND SYSTEMS

Organs are composed of two or more primary tissues that serve the different functions of the organ. The skin is an

organ that has numerous functions provided by its constituent tissues.

LEARNING OUTCOMES

After studying this section, you should be able to:

- 8. Use the skin as an example to describe how the different primary tissues compose organs.
- 9. Identify the body fluid compartments.

An **organ** is a structure composed of at least two, and usually all four, primary tissues. The largest organ in the body, in terms of surface area, is the skin (fig. 1.22). In this section, the numerous functions of the skin serve to illustrate how primary tissues cooperate in the service of organ physiology.

An Example of an Organ: The Skin

The cornified *epidermis* protects the skin against water loss and against invasion by disease-causing organisms. Invaginations of the epithelium into the underlying connective tissue *dermis* create the exocrine glands of the skin. These include hair follicles (which produce the hair), sweat glands, and sebaceous glands. The secretion of sweat glands cools the body by evaporation and produces odors that, at least in lower animals, serve as sexual attractants. Sebaceous glands secrete oily sebum into hair follicles, which transport the sebum to the surface of the skin. Sebum lubricates the cornified surface of the skin, helping to prevent it from drying and cracking.

The skin is nourished by blood vessels within the dermis. In addition to blood vessels, the dermis contains wandering white blood cells and other types of cells that protect against invading disease-causing organisms. It also contains nerve fibers and adipose (fat) cells; however, most of the adipose cells are grouped together to form the *hypodermis* (a layer beneath the dermis). Although adipose cells are a type of connective tissue, masses of fat deposits throughout the body—such as subcutaneous fat—are referred to as *adipose tissue*.

Sensory nerve endings within the dermis mediate the cutaneous sensations of touch, pressure, heat, cold, and pain. Motor nerve fibers in the skin stimulate effector organs, resulting in, for example, the secretions of exocrine glands and contractions of the arrector pili muscles, which attach to hair follicles and surrounding connective tissue (producing goose bumps). The degree of constriction or dilation of cutaneous blood vessels and therefore the rate of blood flow—is also regulated by motor nerve fibers.

The epidermis itself is a dynamic structure that can respond to environmental stimuli. The rate of its cell division—and consequently the thickness of the cornified layer—increases under the stimulus of constant abrasion. This produces calluses. The skin also protects itself against the dangers of ultraviolet light by increasing its production of *melanin* pigment, which absorbs ultraviolet light while producing a tan. In addition, the skin is an endocrine gland; it synthesizes and secretes vitamin D




Figure 1.22 A diagram of the skin. The skin is an organ that contains all four types of primary tissues. APIR

(derived from cholesterol under the influence of ultraviolet light), which functions as a hormone.

The architecture of most organs is similar to that of the skin. Most are covered by an epithelium that lies immediately over a connective tissue layer. The connective tissue contains blood vessels, nerve endings, scattered cells for fighting infection, and possibly glandular tissue as well. If the organ is hollow—as with the digestive tract or blood vessels—the lumen is also lined with an epithelium overlying a connective tissue layer. The presence, type, and distribution of muscle tissue and nervous tissue vary in different organs.

Stem Cells

The different tissues of an organ are composed of cells that are highly specialized, or *differentiated*. The process of differentiation begins during embryonic development, when the fertilized egg, or *zygote*, divides to produce three embryonic tissue layers, or *germ layers: ectoderm, mesoderm*, and *endoderm* (chapter 20; see fig. 20.45*a*). During the course of embryonic and fetal development, the three germ layers give rise to the four primary tissues and their subtypes.

The zygote is totipotent-it can produce all of the different specialized cells of the body. After a number of cell division, when the embryo is at the stage when it implants into the mother's uterus, its cells are *pluripotent*—they can produce all of the body cells except those that contribute to the placenta, and so have been called embryonic stem cells. As development proceeds through successive cell divisions, the cells become increasingly differentiated and lose the ability to form unrelated cell types. However, genes are not lost during differentiation. This was first demonstrated in the 1960s by a British scientist, who found that he could produce a fully formed frog by transplanting the nucleus from a differentiated frog intestinal cell into an egg cell's cytoplasm. In 2006 and later, Japanese scientists found that they could transform differentiated fibroblasts into a pluripotent state, similar to embryonic stem cells, by treating the fibroblasts with a few specific regulatory molecules. The 2012 Nobel Prize in Physiology or Medicine was awarded for these discoveries.

Because the specialized cells have a limited lifespan, many organs retain small populations of cells that are less differentiated and more able to divide to become the specialized cell types within the organ. These less-differentiated cells are known as **adult stem cells.** In the bone marrow, for example, the stem cell population gives rise to all of the different blood cells—red blood cells, white blood cells, and platelets (chapter 13). Similarly, there are stem cells in the brain (chapter 8), skeletal muscles (chapter 12), and intestine (chapter 18).

Scientists have recently discovered that there are also stem cells in the bulge region of the hair follicle (fig. 1.23). These stem cells form keratinocytes, which migrate down to the matrix of the hair follicle and divide to form the hair shaft and root sheath. Other stem cells in the region of the hair follicle just above the bulge form new sebaceous gland cells, which have a high turnover. Skin wounds stimulate the migration of stem cells from the hair follicles into the skin between follicles to promote healing of the wounded skin.

The bulge region also contains melanocyte stem cells, which migrate to the matrix of the follicle and give the hair its color. Scientists have now shown that graying of the hair with age is caused by loss of the melanocyte stem cells in the bulge of the hair follicles. The melanocyte stem cells appeared to be present in most of the hair follicles of people aged 20 to 30 and absent from most hair follicles of people aged 70 to 90.

As demonstrated by the stem cells in the bulge of the hair follicle, adult stem cells can form a variety of related cell types; the adult stem cells are therefore described as *multipotent*. The topics of embryonic and adult stem cells are discussed in more detail in the context of embryonic development (chapter 20, section 20.6).

Systems

Organs that are located in different regions of the body and that perform related functions are grouped into **systems.** These include the integumentary system, nervous system, endocrine system, skeletal system, muscular system, circulatory system,




Figure 1.23 The bulge region of the hair follicle with stem cells. Stem cells in this region migrate to form the differentiated cells of the hair follicle, sebaceous gland, and epidermis. APR

immune system, respiratory system, urinary system, digestive system, and reproductive system (table 1.4). By means of numerous regulatory mechanisms, these systems work together to maintain the life and health of the entire organism.

Body-Fluid Compartments

Tissues, organs, and systems can all be divided into two major parts, or compartments. The **intracellular compartment** is that part inside the cells; the **extracellular compartment** is that part outside the cells. Both compartments consist primarily of water—they are said to be *aqueous*. About 65% of the total body water is in the intracellular compartment, while about 35% is in the extracellular compartment. The two compartments are separated by the cell membrane surrounding each cell (chapter 3, section 3.1).

The extracellular compartment is subdivided into two parts. One part is the *blood plasma*, the fluid portion of the blood. The other is the fluid that bathes the cells within the organs of the body. This is called *tissue fluid*, or *interstitial fluid*. In most parts

Table 1.4 | Organ Systems of the Body

System	Major Organs	Primary Functions
Integumentary	Skin, hair, nails	Protection, thermoregulation
Nervous	Brain, spinal cord, nerves	Regulation of other body systems
Endocrine	Hormone-secreting glands, such as the pituitary, thyroid, and adrenals	Secretion of regulatory molecules called hormones
Skeletal	Bones, cartilages	Movement and support
Muscular	Skeletal muscles	Movements of the skeleton
Circulatory	Heart, blood vessels, lymphatic vessels	Movement of blood and lymph
Immune	Bone marrow, lymphoid organs	Defense of the body against invading pathogens
Respiratory	Lungs, airways	Gas exchange
Urinary	Kidneys, ureters, urethra	Regulation of blood volume and composition
Digestive	Mouth, stomach, intestine, liver, gall- bladder, pancreas	Breakdown of food into molecules that enter the body
Reproductive	Gonads, external genitalia, associated glands and ducts	Continuation of the human species

of the body, blood plasma and tissue fluid communicate freely through blood capillaries. The kidneys regulate the volume and composition of the blood plasma, and thus, indirectly, the fluid volume and composition of the entire extracellular compartment.

There is also selective communication between the intracellular and extracellular compartments through the movement of molecules and ions through the cell membrane, as described in chapter 6. This is how cells obtain the molecules they need for life and how they eliminate waste products.

CHECKPOINTS

- 8a. State the location of each type of primary tissue in the skin.
- **8b.** Describe the functions of nervous, muscle, and connective tissue in the skin.
- **8c.** Describe the functions of the epidermis and explain why this tissue is called "dynamic."
- **9.** Distinguish between the intracellular and extracellular compartments and explain their significance.

Clinical Investigation SUMMARY

Homeostasis of body temperature and blood glucose is maintained by physiological mechanisms that operate in negative feedback loops. Health requires that the body maintain homeostasis, and this can be assessed clinically by specific measurements. If Linda's fasting blood glucose were above the normal range, it could indicate a disease called diabetes mellitus. In this disease, the hormone insulin does not lower the blood glucose concentration adequately in response to a rise in blood glucose.

You can solve another medical mystery now by going to the Connect site for this text. Look for Feedback Control of Blood Glucose, which is based on chapter 1 material. The more you play with physiological concepts, the better you will understand them. The Clinical Investigation boxes at the end of subsequent chapters will provide additional Clinical Investigations on the website that relate to those chapters.

SUMMARY

1.1 Introduction to Physiology 2

- **A.** Physiology is the study of how cells, tissues, and organs function.
 - **1.** In the study of physiology, cause-and-effect sequences are emphasized.
 - **2.** Knowledge of physiological mechanisms is deduced from data obtained experimentally.
- **B.** The science of physiology shares knowledge with the related sciences of pathophysiology and comparative physiology.
 - 1. Pathophysiology is concerned with the functions of diseased or injured body systems and is based on knowledge of how normal systems function, which is the focus of physiology.
 - **2.** Comparative physiology is concerned with the physiology of animals other than humans and shares much information with human physiology.
- **C.** All of the information in this book has been gained by applications of the scientific method. This method has three essential characteristics:
 - **1.** It is assumed that the subject under study can ultimately be explained in terms we can understand.
 - **2.** Descriptions and explanations are honestly based on observations of the natural world and can be changed as warranted by new observations.
 - **3.** Humility is an important characteristic of the scientific method; the scientist must be willing to change his or her theories when warranted by the weight of the evidence.

1.2 Homeostasis and Feedback Control 4

- **A.** Homeostasis refers to the dynamic constancy of the internal environment.
 - **1.** Homeostasis is maintained by mechanisms that act through negative feedback loops.
 - a. A negative feedback loop requires (1) a sensor that can detect a change in the internal environment and (2) an effector that can be activated by the sensor.
 - **b.** In a negative feedback loop, the effector acts to cause changes in the internal environment that compensate for the initial deviations that were detected by the sensor.
 - **2.** Positive feedback loops serve to amplify changes and may be part of the action of an overall negative feedback mechanism.
 - **3.** The nervous and endocrine systems provide extrinsic regulation of other body systems and act to maintain homeostasis.
 - **4.** The secretion of hormones is stimulated by specific chemicals and is inhibited by negative feedback mechanisms.
- **B.** Effectors act antagonistically to defend the set point against deviations in any direction.

1.3 The Primary Tissues 10

- **A.** The body is composed of four types of primary tissues: muscle, nervous, epithelial, and connective tissues.
 - **1.** There are three types of muscle tissue: skeletal, cardiac, and smooth muscle.
 - **a.** Skeletal and cardiac muscle are striated.
 - **b.** Smooth muscle is found in the walls of the internal organs.
 - **2.** Nervous tissue is composed of neurons and neuroglial cells.
 - **a.** Neurons are specialized for the generation and conduction of electrical impulses.
 - **b.** Neuroglial cells provide the neurons with anatomical and functional support.
 - 3. Epithelial tissue includes membranes and glands.
 - **a.** Epithelial membranes cover and line the body surfaces, and their cells are tightly joined by junctional complexes.
 - **b.** Epithelial membranes may be simple or stratified, and their cells may be squamous, cuboidal, or columnar.
 - **c.** Exocrine glands, which secrete into ducts, and endocrine glands, which lack ducts and secrete hormones into the blood, are derived from epithelial membranes.
 - **4.** Connective tissue is characterized by large intercel lular spaces that contain extracellular material.
 - **a.** Connective tissue proper is categorized into subtypes, including loose, dense fibrous, adipose, and others.
 - **b.** Cartilage, bone, and blood are classified as connective tissues because their cells are widely spaced with abundant extracellular material between them.

1.4 Organs and Systems 18

- **A.** Organs are units of structure and function that are composed of at least two, and usually all four, of the primary types of tissues.
 - **1.** The skin is a good example of an organ.
 - **a.** The epidermis is a stratified squamous keratinized epithelium that protects underlying structures and produces vitamin D.
 - **b.** The dermis is an example of loose connective tissue.
 - **c.** Hair follicles, sweat glands, and sebaceous glands are exocrine glands located within the dermis.
 - **d.** Sensory and motor nerve fibers enter the spaces within the dermis to innervate sensory organs and smooth muscles.
 - **e.** The arrector pili muscles that attach to the hair follicles are composed of smooth muscle.
 - 2. Organs that are located in different regions of the body and that perform related functions are grouped into systems. These include, among others, the circulatory system, digestive system, and endocrine system.
 - **3.** Many organs contain adult stem cells, which are able to differentiate into a number of related cell types.
 - **a.** Because of their limited flexibility, adult stem cells are described as multipotent, rather than as totipotent or pluripotent.
 - **b.** For example, the bulge region of a hair follicle contains stem cells that can become keratinocytes, epithelial cells, and melanocytes; the loss of the melanocyte stem cells causes graying of the hair.
- **B.** The fluids of the body are divided into two major compartments.
 - **1.** The intracellular compartment refers to the fluid within cells.
 - **2.** The extracellular compartment refers to the fluid outside of cells; extracellular fluid is subdivided into plasma (the fluid portion of the blood) and tissue (interstitial) fluid.

REVIEW ACTIVITIES

Test Your Knowledge

1. Glands are derived from

b. connective tissue.

- a. nervous tissue.
- d. epithelial tissue.

c. muscle tissue.

- 2. Cells joined tightly together are characteristic of
 - a. nervous tissue.
- c. muscle tissue.
- b. connective tissue. d. epithelial tissue.
- 3. Cells are separated by large extracellular spaces in
 - a. nervous tissue.
- c. muscle tissue.d. epithelial tissue.
- 4. Blood vessels and nerves are usually located within
 - a. nervous tissue.
 - b. connective tissue.

b. connective tissue.

- c. muscle tissue.
- d. epithelial tissue.

- 5. Most organs are composed of
 - a. epithelial tissue.
 - b. muscle tissue.
 - c. connective tissue.
 - d. all of these.
- 6. Sweat is secreted by exocrine glands. This means that
 - a. it is produced by endocrine cells.
 - b. it is a hormone.
 - c. it is secreted into a duct.
 - d. it is produced outside the body.
- 7. Which of these statements about homeostasis is *true*?
 - a. The internal environment is maintained absolutely constant.
 - b. Negative feedback mechanisms act to correct deviations from a normal range within the internal environment.
 - c. Homeostasis is maintained by turning effectors on and off.
 - d. All of these are true.