







Enhanced  
**DIGITAL  
VERSION**  
Included.  
*Details inside.*

**Fred A. Mettler, Jr.**

# ESSENTIALS OF RADIOLOGY

**FOURTH EDITION**



ELSEVIER



# Essentials of Radiology



# Essentials of Radiology

FOURTH EDITION

**Fred A. Mettler, Jr., MD, MPH**

Emeritus Professor  
Department of Radiology  
University of New Mexico  
School of Medicine  
Health Sciences Center  
Albuquerque, New Mexico

ELSEVIER

# ELSEVIER

1600 John F. Kennedy Blvd.  
Ste 1600  
Philadelphia, PA 19103-2899

ESSENTIALS OF RADIOLOGY: FOURTH EDITION  
Copyright © 2019 by Elsevier, Inc. All rights reserved.

ISBN: 978-0-323-50887-2

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the Publisher. Details on how to seek permission, further information about the Publisher's permissions policies, and our arrangements with organizations such as the Copyright Clearance Center and the Copyright Licensing Agency can be found at our website: [www.elsevier.com/permissions](http://www.elsevier.com/permissions).

This book and the individual contributions contained in it are protected under copyright by the Publisher (other than as may be noted herein).

## Notices

Knowledge and best practice in this field are constantly changing. As new research and experience broaden our understanding, changes in research methods, professional practices, or medical treatment may become necessary.

Practitioners and researchers must always rely on their own experience and knowledge in evaluating and using any information, methods, compounds, or experiments described herein. In using such information or methods, they should be mindful of their own safety and the safety of others, including parties for whom they have a professional responsibility.

With respect to any drug or pharmaceutical products identified, readers are advised to check the most current information provided (i) on procedures featured or (ii) by the manufacturer of each product to be administered, to verify the recommended dose or formula, the method and duration of administration, and contraindications. It is the responsibility of practitioners, relying on their own experience and knowledge of their patients, to make diagnoses, to determine dosages and the best treatment for each individual patient, and to take all appropriate safety precautions.

To the fullest extent of the law, neither the Publisher nor the authors, contributors, or editors assume any liability for any injury and/or damage to persons or property as a matter of products liability, negligence, or otherwise or from any use or operation of any methods, products, instructions, or ideas contained in the material herein.

Previous editions copyrighted 2014, 2005, 1996 by Saunders, an imprint of Elsevier Inc.

**Library of Congress Control Number: 2018951185**

*Publisher:* Russell Gabbedy  
*Senior Content Development Specialist:* Ann Anderson  
*Publishing Services Manager:* Catherine Albright Jackson  
*Senior Project Manager:* Doug Turner  
*Designer:* Maggie Reid

Printed in China

Last digit is the print number: 9 8 7 6 5 4 3 2 1



# Preface

Radiology continues to get little attention in most medical school curricula, except perhaps as an elective. The classic gross anatomy lab has been dead (so to speak) for decades, and most health care providers learn human internal anatomy through radiology or electronic formats. This text is not meant to make you a radiologist, nor is it simply a book for a medical student elective, although it has been widely used for the latter. It is intended to be a text that will provide you with a basis for radiologic anatomy, imaging fundamentals, and appropriate imaging for most common clinical problems and be useful for years in your practice. The text is generally organized by clinical presentation (e.g., low back pain, headache) and discusses the imaging that is initially appropriate and why. As such, this text has found wide use among medical students, first year radiology residents, primary care physicians, nurse practitioners, physician's assistants, and other health care professionals.

The fourth edition comes 13 years after the publication of the first edition. Why is a fourth edition needed?

There remains rapid transition in the imaging field, with changes in detector systems to solid state, development of new techniques (e.g., breast tomosynthesis), and continuing development of appropriateness criteria. In the last several years almost 100 new criteria have been developed, which are included in this text. Screening guidelines have been changing, and "rules" have been developed to minimize unnecessary examinations and radiation dose. These areas have all been updated in this edition without expanding the length of the text. While appropriate imaging software is included in some hospital image procedure ordering systems, this is neither widespread nor readily available to you on your smart phone or tablet, but this text and its images are.

I hope that this book fits your needs and wish you the best in your career.

**Fred A. Mettler, Jr.**

# Acknowledgments

I thank my colleagues who have helped me with this edition and previous editions, including Blaine Hart, MD; Charles Hickam, MD; Peter Humphrey, MD; and Josh Robertson, MD. I also thank Gary Mlady, MD, and RuthAnne Bump for their encouragement and help. And a particular note of

gratitude goes out to all those who have worked very hard over the years on many task groups to compile information and recommendations for the American College of Radiology Appropriateness Criteria, which have been essential for this text.



# Contents

- 1 Introduction, 1
- 2 Head and Soft Tissues of Face and Neck, 11
- 3 Chest, 36
- 4 Breast, 93
- 5 Cardiovascular System, 98
- 6 Gastrointestinal System, 125
- 7 Genitourinary System and Retroperitoneum, 168
- 8 Skeletal System, 199
- 9 Nonskeletal Pediatric Imaging, 291
- Appendix, 310



# 1

## Introduction

### AN APPROACH TO IMAGE INTERPRETATION

The first step in medical imaging is to examine the patient and determine the possible cause of his or her problem. Only after this is done can you decide which imaging study is the most appropriate. A vast number of algorithms and guidelines have been developed, but no definite consensus exists on the “right” one for a given symptom or disease because a number of imaging modalities have similar sensitivities and specificities. In this text I provide tables of appropriate initial imaging studies for various clinical situations. When possible, these tables are based on the published literature and recommendations of professional societies. When this is not possible, I give you my opinion based on 45 years of clinical practice.

What should you expect from an imaging examination? Typically one expects to find the exact location of a problem and hopes to make the diagnosis. Although some diseases present a characteristic picture, most can appear in a variety of forms, depending on the stage. As a result, image interpretation will yield a differential diagnosis that must be placed in the context of the clinical findings.

Examination of images requires a logical approach. First you must understand the type of image, the orientation, and the limitations of the technique used. For example, I begin by mentally stating, “I am looking at a coronal computed tomography (CT) scan of the head done with intravenous contrast.” This is important, because intravenous contrast can be confused with fresh blood in the brain.

Next I look at the name and age on the image label to avoid mixing up patients, and this allows making a differential diagnosis that applies to a patient of that age and sex. You would not believe the number of times that this seemingly minor step will keep you from making dumb mistakes.

The next step is to determine the abnormal findings on the image. This means that you need to know the normal anatomy and variants of that particular part of the body, as well as their appearance on the imaging technique used. After this, you should describe the abnormal areas, because it will help you mentally order a differential diagnosis. The most common mistake is to look at an abnormal image and immediately name a disease. When you do this, you will

find your mind locked on that diagnosis (often the wrong one). It is better to say to yourself something like, “I am going to give a differential diagnosis of generalized cardiac enlargement with normal pulmonary vasculature in a 40-year-old man,” rather than to blurt out “viral cardiomyopathy” in a patient who really has a malignant pericardial effusion.

After practicing for 20 years or so, a radiologist knows the spots where pathology most commonly is visualized. Throughout this text, I point out the high-yield areas for the different examinations. Although no absolute rules exist, knowing the pathology and natural history of different diseases will help you. For example, colon cancer typically metastasizes first to the liver rather than the lungs, whereas sarcomas preferentially metastasize to the lungs rather than the liver.

After reviewing the common causes of the imaging findings that you have observed, you should reorder the causes in light of the clinical findings. At this point, you probably think that you are finished. Not so. Often a plethora of information is contained in the patient’s image files or in the hospital’s computer information system. This comes in the form of previous findings and histories supplied for the patient’s other imaging examinations. Reviewing the old reports has directed me to areas of pathology on the current image that I would have missed if I had not looked into the medical information system. A simple example is a pneumonia that has almost but not completely resolved or a pulmonary nodule that, because of inspiratory difference, is hiding behind a rib on the current examination.

You probably think that you are finished now. Wrong again. A certain number of entities could cause the findings on the image, but you just have not thought of them all. After I have finished looking at a case, I try to go through a set sequence of categories in search of other differential possibilities. The categories I use are congenital, physical/chemical, infectious, neoplastic, metabolic, circulatory, and miscellaneous.

### X-RAY

Regular x-rays (plain x-rays, also sometimes called *radiographs*) account for about 75% of imaging examinations. X-ray examinations, or plain x-rays, are made by an x-ray

beam passing through the patient. The x-rays are absorbed in different amounts by the various tissues or materials in the body. Most of the beam is absorbed or scattered. This represents deposition of energy in the tissue but does not cause the patient to become radioactive or to emit radiation. A small percentage of the incident radiation beam exits the patient and strikes a detector.

The historical imaging receptor was a film/screen combination. The x-ray beam would strike a fluorescent screen, which produced light that exposed the film, and then the film was developed. Newer systems are called *computed radiography* or *digital radiography*. In computed radiography, the x-rays strike a plate that absorbs the x-rays and stores the energy at a specific location. The plate is then scanned by a laser, which releases a point of light from the plate. The location is detected and stored in a computer. In digital radiography detector systems, the x-ray hits a detector and then is converted to light or an electrical charge immediately. Once either type of image is stored in the computer, it can be displayed on a monitor for interpretation or transmitted to remote locations for viewing.

Four basic tissue densities, or shades, are visible on plain x-rays. These are air, fat, water (blood and soft tissue), and bone. Air is black or very dark. On regular x-rays and CT scans, fat is generally gray and darker than muscle or blood (Fig. 1.1). Bone and calcium appear almost white. Items that contain metal (such as prosthetic hips) and contrast agents also appear white. The contrast agents generally used are barium for most gastrointestinal studies and iodine for most intravenously administered agents.


Remember that standard or plain x-rays are two-dimensional presentations of three-dimensional information. That is why frontal and lateral views are often needed. Without these, mistakes can easily be made. You must remember that an object visualized on a specific view is somewhere in the path of the x-ray beam (not necessarily in the patient). If an object projects outside the patient on any view, it is outside the patient. However, even if an object

projects within the patient on two orthogonal views, it can still be located outside the patient (Figs. 1.2 and 1.3). Each additional view needed to make a diagnosis requires an additional x-ray exposure and therefore adds to the patient's radiation dose. Radiation doses from various examinations are given in the Appendix.


The terminology used to describe images is usually quite straightforward. Chest and abdominal radiographs are referred to as upright or supine, depending on the position of the patient. In addition, chest x-rays are usually described as posteroanterior (PA) or anteroposterior (AP) (Fig. 1.4). These terms indicate the direction in which the x-ray beam traversed the patient on its way to the detector. PA means that the x-ray beam entered the posterior aspect of the patient and exited anteriorly. AP means that the beam direction through the patient was anterior to posterior. A left lateral decubitus view is one taken with the patient's left side down.

Position is important to note, because it can affect magnification, organ position, and blood flow and therefore significantly affect image interpretation. For example, the heart appears larger on AP than on PA images because on an AP projection the heart is farther from the detector and is magnified more by the diverging x-ray beam. It also appears larger on supine than on upright images because the hemidiaphragms are pushed up, making the heart appear wider. Portable chest images are taken not only in the AP projection but also with the tube closer to the patient than on standard upright images. This magnifies the heart even more.


Use of contrast agents permits visualization of anatomic structures that are not normally seen. For example, intravenously or intra-arterially injected agents allow visualization of blood vessels (Fig. 1.5). If imaging is done with standard format, the blood vessels appear white. Digital imaging allows subtraction or removal of unwanted structures, such as the bones, from an image (see Fig. 1.5B). Often the computer manipulation is done in such a way that the



• **Fig. 1.1** The Four Basic Densities on an X-Ray. A lateral view of the forearm shows that the bones are the densest, or white; soft tissue is gray; fat is somewhat dark; and air is very dark. The abnormality in this case is the fat in the soft tissue of the forearm, which is due to a lipoma.



• **Fig. 1.2** Spatial Localization on an X-Ray. On both anteroposterior (AP) and lateral projections, the square and round objects will be seen projecting within the view of the chest, even though the square object is located outside the chest wall. If you can see an object projecting outside the chest wall on at least one view (the triangle), it is outside the chest. If, however, an object looks as though it is inside the chest on both views, it may be either inside or outside.



• **Fig. 1.3** What Is the Location of the Keys? On both the posteroanterior (PA) view of the chest (A) and the lateral view (B), the keys seem to be within the center of the chest. Actually, if you look carefully, you will notice that the keys do not change position at all, even though the patient has rotated 90 degrees. The keys are located on the receptor cassette and are not in the patient.


arteries may appear black instead of white, although this usually does not present a problem in interpretation.

Contrast agents are used to fill either a hollow viscus (such as the stomach) or anatomic tubular structures that can be accessed in some way (such as blood vessels, ureter, and common bile duct). When you see an abnormality on one of these studies, you must determine whether the location is intraluminal, mural, or extrinsic. This usually requires seeing the abnormality in perpendicular views (Fig. 1.6). Unless you are careful about this determination, you will make errors in diagnosis.

Contrast agents instilled orally, rectally, or retrograde into the ureter or bladder incur little or no risk unless

aspiration or perforation occurs. With the intravenously or intra-arterially administered agents, a small but real risk for contrast reaction exists. This is something that you should consider before ordering a contrast-enhanced CT scan. About 5% of patients will experience an immediate mild reaction, such as a metallic taste or a feeling of warmth; some experience nausea and vomiting, wheeze, or get hives as a result of these contrast agents. Some of these mild reactions can be treated with 50 mg of intramuscular diphenhydramine (Benadryl). Because contrast agents also can reduce renal function, they should not generally be used in patients with compromised renal function (estimated glomerular filtration rate [eGFR] <50 to 60 mL/min).

About 1 in 1000 patients have a severe reaction to intravascular contrast. This may be a vasovagal reaction, laryngeal edema, severe hypotension, an anaphylactic-type reaction, or cardiac arrest. A vasovagal reaction can be treated with 0.5 to 1.0 mg of intravenous atropine. The most important initial therapeutic measures for these severe reactions are to establish an airway, ensure breathing and circulation, and give intravenous fluids. Other drugs



• **Fig. 1.4** Typical X-ray Projections. X-ray projections are typically listed as anteroposterior (AP) or posteroanterior (PA). This depends on whether the x-ray beam passed through the patient from anterior to posterior or the reverse. Lateral (LAT) and oblique (OBL) views also are commonly obtained.


obviously also may be necessary. The risk for death from a study using intravenously administered contrast agents is between 1 in 40,000 and 1 in 100,000.

## COMPUTED TOMOGRAPHY


CT is accomplished by passing a rotating fan beam of x-rays through the patient and measuring the transmission at thousands of points. The data are handled by a computer that calculates exactly what the x-ray absorption was at any given spot in the patient. The data can be manipulated in a number of ways, displayed on a screen, or photographed. Because the data points are in the computer memory, it is possible to “window” the data and obtain a number of images without additional radiation exposure (Fig. 1.7). The computers can even display the data as a three-dimensional rotating image, although this is rarely necessary for diagnosis. Compared with plain x-rays, CT uses about 10 to 100 times more radiation.

On early CT scanners the x-ray tube rotated around the patient to obtain a single “slice,” and then the table was moved incrementally before another slice was obtained. Newer scanners allow the x-ray tube to stay on and rotate at the same time that the table is moving. This is called a *spiral scanner* or *helical scanner*. The most modern scanners not only have the helical motion but also have multiple rows of detectors and can obtain more than 100 image data slices at once.


The appearance of tissues on CT scan depends to some extent on the computer manipulation, but in general the



• **Fig. 1.5** Pulmonary Angiogram. (A) A conventional view of blood vessels can be obtained by injecting iodinated contrast material into the vessels. On these images the vessels will appear white and the bones will be seen as you would normally expect (white). A digital subtraction technique with a computer may show the vessels either as black (B) or as white, but the bones will have been subtracted from the image.



• **Fig. 1.6** Appearances of Different Lesions Depending on Their Location When Using Contrast. Contrast medium is used to visualize tubular structures, including the spinal canal, blood vessels, gastrointestinal tract, ureters, and bladder. (A) Intraluminal lesions, such as stones or blood clots within the lumen of the given structure, produce a central defect on both anteroposterior (AP) and lateral projections. On the AP and lateral views the contrast will show acute angles on both sides and in both projections. (B) Intramural lesions will produce a defect that indents the column of contrast. When seen tangentially, an acute angle will appear between the normal wall and the beginning of the indentation. (C) Extramural lesions also can indent the wall, but at the point of indentation, the angle will be somewhat blunted as compared with the intramural lesion.



• **Fig. 1.7** Computed Tomography (CT). Images of the abdomen are presented here. (A) The image was made by using relatively wide windows during viewing, and no intravenous contrast was used. (B) The windows have been narrowed, producing a rather grainy image, and intravenous contrast has been administered so that you can see enhancement of the aorta, abdominal vessels, and both kidneys (K). In both images, contrast has been put in the bowel (B) to differentiate bowel from solid organs and structures. L, Left; R, right; Sp, spine.

basic four densities on CT images are the same as those in plain x-rays: air is black, fat is dark gray, soft tissue is light gray, and bone or calcium and contrast agents are white. One advantage of CT is that actual x-ray absorption of a specific tissue can be displayed. The units used are Hounsfield units. The Hounsfield density of water is zero. The greater sensitivity of CT compared with plain x-rays allows areas of tiny punctate calcification to be seen.

CT scans are presented as a series of slices of tissue. The method is similar in principle to slicing a loaf of bread and pulling up one slice at a time to examine it. Thus CT is a two-dimensional display of two-dimensional information, and objects appear where they really are in space. The scans or slices are shown as if you were viewing the patient from the foot of the patient's bed. Thus the individual's right side is on your left (Fig. 1.8). This also is the convention used for transverse images of ultrasound and magnetic resonance imaging (MRI).

Contrast agents, frequently used in CT scans, are usually the same water-soluble oral, rectal, or intravenous iodinated agents used in other imaging studies. Intravenous contrast agents are common, being used in probably 75% of all CT studies, and obviously carry the risk for contrast reactions discussed previously. Rapid acquisition of images allows the intravenously administered contrast to be displayed and images acquired in arterial, venous, or delayed phases with only a single injection.

The appeal of CT is that a large number of structures are visualized simultaneously. In a patient with abdominal pain, one CT examination shows the liver, adrenal glands, kidneys, spleen, aorta, pancreas, and other structures. This allows the clinician to identify macroscopic pathology quickly.