


ROGER P. SMITH

NETTER'S OBSTETRICS & GYNECOLOGY

3rd EDITION

*F. Netter
M.D.*



ELSEVIER

Any screen. Any time. Anywhere.

Activate the eBook version
of this title at no additional charge.



Expert Consult eBooks give you the power to browse and find content, view enhanced images, share notes and highlights—both online and offline.

Unlock your eBook today.

- 1 Visit expertconsult.inkling.com/redeem
- 2 Scratch off your code
- 3 Type code into “Enter Code” box
- 4 Click “Redeem”
- 5 Log in or Sign up
- 6 Go to “My Library”

It's that easy!

Scan this QR code to redeem your eBook through your mobile device:



Place Peel Off
Sticker Here

For technical assistance:
email expertconsult.help@elsevier.com
call 1-800-401-9962 (inside the US)
call +1-314-447-8200 (outside the US)

ELSEVIER

Use of the current edition of the electronic version of this book (eBook) is subject to the terms of the nontransferable, limited license granted on expertconsult.inkling.com. Access to the eBook is limited to the first individual who redeems the PIN, located on the inside cover of this book, at expertconsult.inkling.com and may not be transferred to another party by resale, lending, or other means.

NETTER:


It's How You Know Anatomy.



Entire Collection Now Available!

Netter Collection of Medical Illustrations, 2nd Edition

The Netter Collection of Medical Illustrations, Dr. Frank H. Netter's decades of work devoted to depicting each of the major body systems, has been updated and brought into modern context. The second edition of the legendary "green books" offers **Netter's timeless work**, now arranged and enhanced by **modern text and radiologic imaging**. **Contributions** by field-leading doctors and teachers from world-renowned medical institutions are supplemented with **new illustrations** created by **master artist-physician Carlos Machado** and other top medical illustrators working in the Netter tradition.




Netter's Correlative Imaging Series

The *Netter's Correlative Imaging* series pairs classic Netter and Netter-style illustrations with imaging studies and succinct descriptions to provide you with a richer understanding of human anatomy. These comprehensive, normal anatomy atlases cover all major sections of the body, featuring illustrated plates side-by-side with the most common imaging modalities for each region.

Shop online at elsevierhealth.com


Stay current and connected in ObGyn with coverage of the latest hot topics.



Comprehensive Gynecology, 7th Edition

Rogério A. Lobo, MD, David M. Gershenson, MD, Gretchen M. Lentz, MD, and Fidel A. Valea, MD

The primary gynecology text for over 25 years, *Comprehensive Gynecology* covers all of the **key issues** residents may encounter. This 7th edition has been **fully updated with a wealth of new content**, including current discussions of minimally invasive surgical approaches to gynecologic care, infertility issues and treatments, effectively managing menopausal patients, and more.



Obstetrics: Normal and Problem Pregnancies, 7th Edition

Steven G. Gabbe, MD, Jennifer R. Niebyl, MD, Joe Leigh Simpson, MD, Mark B. Landon, MD, Henry L. Galan, MD, Eric R.M. Jauniaux, MD, PhD, Deborah A. Driscoll, MD, Vincenzo Berghella, MD, and William A. Grobman, MD



Obstetrics: Normal and Problem Pregnancies remains a go-to **choice for authoritative guidance** on managing today's obstetric patient. **International experts** put the latest knowledge in this specialty at your fingertips, with current and relevant information on everything from fetal origins of adult disease, to improving global maternal health, to important topics in day-to-day obstetrical practice. **Highly readable, well illustrated, and easy to understand**, this bestselling obstetrics reference is an ideal tool for residents.

Shop online at elsevierhealth.com!

ELSEVIER

NETTER'S OBSTETRICS & GYNECOLOGY

3rd EDITION



Roger P. Smith, MD

Assistant Dean for Graduate Medical Education
Professor of Clinical Biologic Sciences
Charles E. Schmidt College of Medicine
Florida Atlantic University
Boca Raton, Florida

Illustrations by Frank H. Netter, MD

Contributing Illustrators

Carlos A.G. Machado, MD

John A. Craig, MD

Kristen Wienandt Marzejon, MS, MFA

Joe Chovan

James A. Perkins, MS, MFA

Tiffany S. DaVanzo, MA, CMI

ELSEVIER

ELSEVIER

1600 John F. Kennedy Blvd.
Ste 1800
Philadelphia, PA 19103-2899

NETTER'S OBSTETRICS AND GYNECOLOGY, THIRD EDITION

ISBN: 978-0-7020-7036-5

Copyright © 2018 by Elsevier, Inc. All rights reserved.
Previous edition copyrighted 2009

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. Details on how to seek permission, further information about the Publisher's permissions policies and our arrangements with organizations such as the Copyright Clearance Center and the Copyright Licensing Agency, can be found at our website: www.elsevier.com/permissions.

This book and the individual contributions contained in it are protected under copyright by the Publisher (other than as may be noted herein).

Notices

Knowledge and best practice in this field are constantly changing. As new research and experience broaden our understanding, changes in research methods, professional practices, or medical treatment may become necessary.

Practitioners and researchers must always rely on their own experience and knowledge in evaluating and using any information, methods, compounds, or experiments described herein. In using such information or methods they should be mindful of their own safety and the safety of others, including parties for whom they have a professional responsibility.

With respect to any drug or pharmaceutical products identified, readers are advised to check the most current information provided (i) on procedures featured or (ii) by the manufacturer of each product to be administered, to verify the recommended dose or formula, the method and duration of administration, and contraindications. It is the responsibility of practitioners, relying on their own experience and knowledge of their patients, to make diagnoses, to determine dosages and the best treatment for each individual patient, and to take all appropriate safety precautions.

To the fullest extent of the law, neither the Publisher nor the authors, contributors, or editors, assume any liability for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions, or ideas contained in the material herein.

Library of Congress Cataloging-in-Publication Data

Names: Smith, Roger P. (Roger Perry), 1949- author. | Netter, Frank H. (Frank Henry), 1906-1991, illustrator.

Title: Netter's obstetrics and gynecology / Roger P. Smith ; illustrations by Frank H. Netter ; contributing illustrators, Carlos A.G. Machado, John A. Craig, Kristen Wienandt Marzejon, Joe Chovan, James A. Perkins, Tiffany S. DaVanzo.

Other titles: Obstetrics and gynecology

Description: Third edition. | Philadelphia, PA : Elsevier, [2017] | Includes bibliographical references and index.

Identifiers: LCCN 2016051375 | ISBN 9780702070365 (hardcover : alk. paper)

Subjects: | MESH: Genital Diseases, Female | Obstetric Surgical Procedures | Pregnancy Complications | Atlases

Classification: LCC RG110 | NLM WP 17 | DDC 618.1-dc23 LC record available at <https://lccn.loc.gov/2016051375>

Content Strategist and Content Development Specialist: Marybeth Thiel
Publishing Services Manager: Patricia Tannian
Project Manager: Ted Rodgers
Design Direction: Patrick Ferguson

Printed in China

Last digit is the print number: 9 8 7 6 5 4 3 2 1



Working together
to grow libraries in
developing countries

www.elsevier.com • www.bookaid.org

PREFACE

No student of medicine, past or present, is unaware of the extraordinary series of medical illustrations created by Dr. Frank Netter. It is an incredible body of work that has been carried forward by the talented Carlos Machado, MD, and John Craig, MD, since Dr. Netter's passing. Older physicians have looked with envy at these images, wishing they had been available when they were learning; established physicians return to them as comfortable sources of information; young physicians seek them out for the wealth of information they contain and their ability to make clear difficult clinical concepts. This spirit of concise reference and resource is the premise of this text.

This third edition maintains the same consistent format in presenting topics to facilitate rapid access—the same information is in the same location—that was so well received in the first and second

editions. Chapters have been organized to provide a quick, concise resource for the diagnosis and treatment of common conditions encountered by anyone who provides care for women. In producing this third edition, more than 25 new topics have been added, including sections on embryology and anatomy, a more intuitive organization has been developed, an expanded section on commonly encountered procedures is included, new artwork has been developed, and subtle enhancements (such as indications of the level of evidence provided for references) have been made throughout the work.

It is our hope that this work will be both a useful resource and a celebration of the artistic richness that is clinical medicine.

Roger P. Smith, MD

ABOUT THE ARTISTS

FRANK H. NETTER, MD

Frank H. Netter was born in 1906 in New York City. He studied art at the Art Students League and the National Academy of Design before entering medical school at New York University, where he received his MD degree in 1931. During his student years, Dr. Netter's notebook sketches attracted the attention of the medical faculty and other physicians, allowing him to augment his income by illustrating articles and textbooks. He continued illustrating as a sideline after establishing a surgical practice in 1933, but he ultimately opted to give up his practice in favor of a full-time commitment to art. After service in the United States Army during World War II, Dr. Netter began his long collaboration with the CIBA Pharmaceutical Company (now Novartis Pharmaceuticals). This 45-year partnership resulted in the production of the extraordinary collection of medical art so familiar to physicians and other medical professionals worldwide.

In 2005, Elsevier purchased the Netter Collection and all publications from Icon Learning Systems. There are now over 50 publications featuring the art of Dr. Netter available through Elsevier, Inc. (in the US: www.us.elsevierhealth.com/Netter; outside the US: www.elsevierhealth.com).

Dr. Netter's works are among the finest examples of the use of illustration in the teaching of medical concepts. The 13-book *Netter Collection of Medical Illustrations*, which includes the greater part of the more than 20,000 paintings created by Dr. Netter, became and remains one of the most famous medical works ever published. The *Netter Atlas of Human Anatomy*, first published in 1989, presents the anatomical paintings from the Netter Collection. Now translated into 16 languages, it is the anatomy atlas of choice among medical and health professions students the world over.

The Netter illustrations are appreciated not only for their aesthetic qualities, but, more important, for their intellectual content. As Dr. Netter wrote in 1949, "... clarification of a subject is the aim and goal of illustration. No matter how beautifully painted, how delicately and subtly rendered a subject may be, it is of little value as a *medical illustration* if it does not serve to make clear some medical point." Dr. Netter's planning, conception, point of view, and approach are what inform his paintings and what make them so intellectually valuable.

Frank H. Netter, MD, physician and artist, died in 1991.

Learn more about the physician-artist whose work has inspired the Netter Reference collection:

<http://www.netterimages.com/artist/netter.htm>.

CARLOS MACHADO, MD

Carlos Machado was chosen by Novartis to be Dr. Netter's successor. He continues to be the main artist who contributes to the Netter collection of medical illustrations.

Self-taught in medical illustration, cardiologist Carlos Machado has contributed meticulous updates to some of Dr. Netter's original plates and has created many paintings of his own in the style of Netter as an extension of the Netter collection. Dr. Machado's photorealistic expertise and his keen insight into the physician/patient relationship inform his vivid and unforgettable visual style. His dedication to researching each topic and subject he paints places him among the premier medical illustrators at work today.

Learn more about his background and see more of his art at:

<http://www.netterimages.com/artist/machado.htm>.

ONLINE CONTENTS

Visit www.ExpertConsult.com for the following:

Printable Patient Education Brochures from Ferri's Netter Patient Advisor, 3rd Edition

1. Managing Your Anorexia Nervosa
2. Breast Cancer: Screening and Prevention
3. Managing Your Bulimia
4. Managing Your Cervical Cancer
5. Managing Your Cervical Incompetence
6. Managing Your Cervical Polyps
7. Managing Your Dysmenorrhea
8. Managing Your Dysfunctional Uterine Bleeding
9. Managing Your Endometriosis
10. Managing Your Female Sexual Dysfunction
11. Managing Your Fibrocystic Breast Disease
12. Managing Your Hyperemesis Gravidarum
13. Managing Your Infertility
14. Kegel Exercises Strengthening Your Pelvic Floor Muscles
15. Managing Your Lactational Mastitis
16. Managing Your Menopause
17. Managing Your Menorrhagia
18. Managing Your Morning Sickness
19. Managing Your Ovarian Cancer
20. Managing Your Ovarian Cysts
21. Managing Your Pelvic Inflammatory Disease
22. Managing Your Uterine Fibroids
23. Managing Your Vaginal Dryness
24. Managing Your Bacterial Vaginal Infection
25. Managing Your Miscarriage
26. Weight Control For Women

CONTENTS

PART I: ANATOMY AND EMBRYOLOGY

SECTION I

Embryology

1 Sexual Differentiation	3
2 Upper Genital Tract Development	4
3 Lower Genital Tract Development	5
4 Development of the Breast	7

SECTION II

Anatomy

5 External Genitalia	11
6 Perineum	12
7 Vagina	13
8 Pelvic Viscera	15
9 Cervix, Uterus, and Adnexa	16
10 Ovaries	17
11 Breast	19

PART II: GYNECOLOGY AND WOMEN'S HEALTH

SECTION III

General Health Considerations and Counseling

12 Puberty: Normal Sequence	25
13 Health Maintenance: Ages 12–18 Years	26
14 Health Maintenance: Ages 19–39 Years	28
15 Contraception: Counseling Principles	30
16 Health Maintenance: Ages 40–64 Years	32
17 Health Maintenance: Ages 65 Years and Older	35

SECTION IV

Diseases, Disorders, and Common Problems

18 Abortion	41
19 Abuse: Physical and Sexual	44
20 Acne	46
21 Alzheimer Disease	48
22 Anemia	50
23 Anorectal Fistula	52
24 Anxiety	54
25 Asthma	56
26 Cholelithiasis	58

27 Chronic Pelvic Pain	60
28 Constipation	62
29 Crohn Disease	64
30 Depression (Unipolar)	66
31 Diverticular Disease	68
32 Domestic Violence	69
33 Dysmenorrhea: Primary and Secondary	71
34 Dyspareunia: Deep Thrust	73
35 Dysuria	75
36 Eating Disorders: Anorexia Nervosa and Bulimia	77
37 Fibromyalgia	79
38 Gallbladder Disease	82
39 Gastritis	84
40 Gastroesophageal Reflux	85
41 Hair Loss	87
42 Headache: Tension and Cluster	89
43 Headache: Migraine	92
44 Hematuria	94
45 Hemorrhoids	96
46 Hyperthyroidism	97
47 Hypothyroidism	99
48 Infertility: General Considerations	101
49 Irritable Bowel Syndrome	103
50 Low Back Pain	105
51 Melanoma	108
52 Myofascial Syndromes	110
53 Obesity	112
54 Osteoporosis	114
55 Pessary Therapy	117
56 Postcoital Bleeding	119
57 Premenstrual Syndrome	121
58 Pruritus Ani	123
59 Rape and Rape Trauma Syndrome	124
60 Sexual Dysfunction: Libidinal and Orgasmic Dysfunction (Anorgasmia)	127
61 Sexually Transmitted Infections: Chancroid	129
62 Sexually Transmitted Infections: <i>Chlamydia trachomatis</i>	131
63 Sexually Transmitted Infections: Condylomata Acuminata	133

64 Sexually Transmitted Infections: Gonorrhea	135	96 Vulvar Hematoma	201
65 Sexually Transmitted Infections: Granuloma Inguinale (Donovanosis)	137	97 Vulvar Lesions	203
66 Sexually Transmitted Infections: Herpes	139	98 Vulvar Vestibulitis (Provoked Vulvodynia)	206
67 Sexually Transmitted Infections: Human Immunodeficiency Virus	141	SECTION VI	
68 Sexually Transmitted Infections: Human Papillomavirus	144	Vaginal Disease	
69 Sexually Transmitted Infections: Lymphogranuloma Venereum	147	99 Cystocele/Urethrocele	211
70 Sexually Transmitted Infections: Molluscum Contagiosum	149	100 Enterocele	212
71 Sexually Transmitted Infections: Parasites	150	101 Fistulae: Gastrointestinal and Urinary Tract	214
72 Sexually Transmitted Infections: Syphilis	152	102 Rectocele	216
73 Sexually Transmitted Infections: <i>Trichomonas vaginalis</i>	155	103 Sarcoma Botryoides	218
74 Thrombophlebitis	157	104 Transverse Vaginal Septum	219
75 Toxic Shock Syndrome	159	105 Vaginal Cysts	221
76 Ulcerative Colitis	162	106 Vaginal Dryness	222
77 Urinary Incontinence: Bypass and Overflow	164	107 Vaginal Lacerations	223
78 Urinary Incontinence: Stress	166	108 Vaginal Prolapse	225
79 Urinary Incontinence: Urge	169	109 Vaginitis: Atrophic	227
80 Urinary Tract Infection	171	110 Vaginitis: Bacterial (Nonspecific) and Bacterial Vaginosis	229
81 Urodynamics Testing	174	111 Vaginitis: Monilial	231
82 Varicose Veins	176	112 Vaginitis: <i>Trichomonas</i>	233
SECTION V		SECTION VII	
Vulvar Disease		Cervical Disease	
83 Acne Inversa (Hidradenitis Suppurativa)	181	113 Abnormal Pap Smear: Atypical Squamous or Glandular Cells of Undetermined Origin	237
84 Bartholin Gland: Abscess/Infection	182	114 Abnormal Pap Smear: Low-Grade Squamous Intraepithelial Lesions and High-Grade Squamous Intraepithelial Lesions	239
85 Bartholin Gland: Cysts	184	115 Carcinoma in Situ (Cervix)	241
86 Contact Vulvitis	186	116 Cervical Cancer	243
87 Dyspareunia: Insertional	187	117 Cervical Erosion	245
88 Female Circumcision	188	118 Cervical Eversion	246
89 Hymenal Stenosis	190	119 Cervical Polyps	247
90 Hyperplastic Vulvar Dystrophy (Squamous Cell Hyperplasia, Lichen Simplex Chronicus)	192	120 Cervical Stenosis	248
91 Imperforate Hymen	193	121 Cervicitis	250
92 Labial Adhesions	194	122 Nabothian Cysts	252
93 Lichen Planus	196	SECTION VIII	
94 Lichen Sclerosus	197	Uterine Pathology	
95 Vulvar Cancer	199	123 Adenomyosis	255
		124 Asherman Syndrome (Uterine Synechia)	256
		125 Dysfunctional (Abnormal) Uterine Bleeding	258
		126 Endometrial Cancer	261
		127 Endometrial Hyperplasia: Simple and Complex	263

128	Endometrial Polyps	265
129	Endometritis	266
130	Hematometra	268
131	Intermenstrual Bleeding	269
132	Irregular Menstrual Periods	271
133	Menorrhagia	274
134	Postmenopausal Vaginal Bleeding	276
135	Sarcoma (Uterine)	278
136	Uterine Anomalies: Bicornuate, Septate, and Unicornuate Uterus	280
137	Uterine Leiomyomata (Fibroids, Myoma)	282
138	Uterine Prolapse	284

SECTION IX

Adnexal Disease

139	Adenofibroma	289
140	Clear Cell Carcinoma	290
141	Dermoid Cyst	292
142	Dysgerminoma	293
143	Ectopic Pregnancy	295
144	Endometriosis	298
145	Epithelial Stromal Ovarian Tumors	301
146	Germ Cell Tumor	303
147	Granulosa Cell Tumors	304
148	Hydrosalpinx (Chronic Pelvic Inflammatory Disease)	307
149	Krukenberg Tumor	309
150	Mucinous Ovarian Cysts	310
151	Ovarian Cancer	312
152	Ovarian Cysts	315
153	Ovarian Fibroma	317
154	Ovarian Torsion	319
155	Pelvic Inflammatory Disease (PID)	321
156	Pseudomyxoma Peritonei	324
157	Serous Ovarian Cysts	325
158	Sertoli-Leydig Cell Tumor (Arrhenoblastoma)	327
159	Transitional Cell (Brenner) Tumor	329

SECTION X

Breast Diseases and Conditions

160	Accessory Nipples (Polythelia)	333
161	<i>BRCA1</i> and <i>BRCA2</i> Mutations	334
162	Breast Cancer	336
163	Breast Cyst	339
164	Breast Duct Ectasia	340
165	Breast Fat Necrosis	342
166	Breast Fibroadenoma	343
167	Breast Intraductal Papilloma	345

168	Fibrocystic Breast Change	347
169	Galactocele	349
170	Galactorrhea	351
171	Mammography	353
172	Mastitis (Lactational)	355
173	Mastodynia and Mastalgia (Breast Pain)	356
174	Mondor Disease	358
175	Nipple Discharge	360
176	Paget Disease of the Breast	362

SECTION XI

Reproductive, Genetic, and Endocrine Conditions

177	Abnormal Puberty	367
178	Amenorrhea: Primary	369
179	Amenorrhea: Secondary	371
180	Androgen Insensitivity Syndrome	372
181	Anovulation	374
182	Assisted Reproduction	376
183	Down Syndrome	379
184	Gonadal Dysgenesis	381
185	Hirsutism	384
186	Hyperprolactinemia	387
187	Infertility	390
188	Menopause	392
189	Polycystic Ovary Syndrome	395
190	Recurrent Pregnancy Loss	397
191	Sexual Ambiguity	399
192	Sheehan Syndrome	401
193	Turner Syndrome	403
194	Uterine Agenesis	405
195	Vaginal Agenesis	407
196	Virilization	408

PART III: OBSTETRICS

SECTION XII

Obstetrics: General Considerations

197	Preconceptional Care and Counseling	413
198	Routine Prenatal Care: First Trimester	415
199	Routine Prenatal Care: Second Trimester	416
200	Routine Prenatal Care: Third Trimester	418
201	Antepartum Fetal Testing	420
202	Biophysical Profile	422

203 Contraction Stress Testing	424	243 Shoulder Dystocia	503
204 Doppler Flow Studies	425	244 Third Trimester Bleeding	505
205 Nonstress Testing	427	245 Trauma in Pregnancy	506
206 Normal Labor	429	246 Uterine Atony and Postpartum Hemorrhage	508
207 Fetal Heart Rate Testing: Bradycardia	431	247 Uterine Inversion	511
208 Fetal Heart Rate Testing: Periodic Changes	433	248 Uterine Rupture	513
209 Fetal Heart Rate Testing: Reduced Variability	435		
210 Fetal Heart Rate Testing: Tachycardia	437		
211 Obstetric Anesthesia/Analgesia	438		
212 Normal Postpartum Changes	440		
213 Breastfeeding (Lactation)	442		
SECTION XIII Obstetric Conditions and Concerns		PART IV: PROCEDURES	
214 Abnormalities of Placental Implantation	447	SECTION XIV Procedures	
215 Active Management of Labor	449	249 Amniocentesis	517
216 Acute Fatty Liver of Pregnancy	450	250 Aspiration of Breast Cyst	518
217 Amniotic Fluid Embolism	452	251 Bartholin Gland Cyst/Abscess Drainage	520
218 Breech Birth	454	252 Bartholin Gland Marsupialization	521
219 Caput Succedaneum	456	253 Breast Biopsy: Core	523
220 Cardiovascular Disease in Pregnancy	457	254 Breast Biopsy: Open	524
221 Cervical Insufficiency	460	255 Cervical Cerclage	526
222 Cholecystitis in Pregnancy	461	256 Cervical Conization (Cold Knife)	528
223 Chorioamnionitis	464	257 Cervical Polypectomy	529
224 Diabetes Mellitus in Pregnancy	466	258 Cesarean Delivery	531
225 Fetal Alcohol Syndrome	468	259 Chorionic Villus Sampling	534
226 Gestational Trophoblastic Disease	470	260 Circumcision (Male; Newborn and Infant)	536
227 Gingivitis in Pregnancy	472	261 Colposcopy	539
228 HELLP Syndrome	474	262 Cervical Cryocautery	541
229 Hepatitis in Pregnancy	476	263 Cystourethroscopy	542
230 Hyperemesis Gravidarum	478	264 Diaphragm Fitting	544
231 Intrauterine Growth Restriction	479	265 Dilation and Curettage	546
232 Multiple Gestation	482	266 Endometrial Biopsy	548
233 Oligohydramnios	484	267 Forceps-Aided Delivery	550
234 Placenta Previa	486	268 Hysteroscopy: Diagnostic	552
235 Placental Abruption	488	269 Hysteroscopy: Polyp and Leiomyoma Resection	554
236 Polyhydramnios	489	270 Hysteroscopic Sterilization	556
237 Postpartum Breast Engorgement	491	271 Intrauterine Contraceptive Device Insertion	558
238 Postpartum Depression	492	272 Intrauterine Contraceptive Device Removal	560
239 Preeclampsia and Eclampsia	494	273 LEEP (Loop Electrosurgical Excision Procedure) and LLETZ (Large Loop Excision of the Transformation Zone) Conizations	562
240 Pruritic Urticarial Papules and Plaques of Pregnancy (PUPPP)	497	274 Pessary Fitting	564
241 Puerperal Infection	499	275 Sonohysterography	566
242 Rh Incompatibility	501	276 Subdermal Contraceptive Capsule Insertion	568
		277 Subdermal Contraceptive Capsule Removal	570
		278 Transvaginal Ultrasonography	571
		279 Trigger Point Injection	573

280 Urodynamic Testing: Complex	574	282 Vacuum-Assisted Delivery	578
281 Urodynamic Testing: Simple	576	Index	581

Embryology

- 1 Sexual Differentiation
- 2 Upper Genital Tract Development
- 3 Lower Genital Tract Development
- 4 Development of the Breast



This page intentionally left blank

Genetic sex is determined by the complement and function of sex chromosomes (X and Y) that are present at the time of conception. A Y chromosome carrying specific genes is necessary for the development of testes. The testes are responsible for the organization of the sexual duct system into a male configuration and for the suppression of the paramesonephric (Müllerian) system responsible for female anatomic structures. In the absence of a Y chromosome, specifically required genes, or a functioning gonad, the development will be female. General phenotypic development of the female

is viewed as a default event, although new evidence of a more complex process is emerging.

Sexual differentiation genes are located on the Y chromosome, the primary of which is the *SRY* gene, also called the testis-determining factor. The *SRY* gene is found on the short arm of the Y chromosome and influences Sertoli cell differentiation, mesonephric ridge cell development, and male architectural development of the gonad, including blood vessels and other structures of the testes. Several other genes, including those that express




Figure 1.1 Genetics and biology of early reproductive tract development

steroidogenic factor-1, WT1, and DAX1, on other chromosomes, are also necessary for normal testicular development. To date, multiple mutations of the *SRY* gene have been reported and all are associated with sex reversals (female phenotype).

As discussed earlier, genes in other locations are also important for complete male sexual differentiation. *DAX1*, a nuclear hormone receptor, can alter *SRY* activity during development by suppressing genes downstream to *SRY* that would normally induce testicular differentiation. A second gene, *WNT4*, largely confined to the adult ovary, may also serve as an “antitestis” gene. In very rare male individuals a Y chromosome may be absent, but the *SRY* gene may be present on another chromosome, most commonly the X chromosome, resulting in a male phenotype. It is becoming apparent that genes, such as *WNT4* and *DAX1*, can proactively induce female gonadal development even in the presence of *SRY*, thereby further complicating the picture. This may account for individuals who are exceptions to the normal sexual dichotomy (eg, males with a uterus or females with an XY karyotype) or who exhibit biologic and/or behavioral characteristics of both sexes.

Male gonadal development precedes female development, and the early secretion of testosterone and anti-Müllerian hormone (AMH) steers the further development of the genital tracts away from the default female phenotype. At a critical point, AMH, produced by Sertoli cells, and testosterone, secreted by Leydig cells, must be produced in sufficient amounts. AMH acts locally, thus

suppressing the Müllerian duct system. Testosterone acts systemically, thus causing the differentiation of the mesonephric duct system and male development of the urogenital tubercle, urogenital sinus, and urogenital folds. Enzymes involved in testosterone biosynthesis and conversion to dihydrotestosterones are regulated by genes located on autosomes. The ability to secrete AMH is a recessive trait coded on either an autosome or the X chromosome, and genes for the development of cytoplasmic receptors of androgens seem to be coded on the X chromosome.

The development of the ovary occurs at approximately the 11th or 12th week of gestation, although the primordial germ cells migrate several weeks earlier to the germinal ridge. Two functional X chromosomes are necessary for the optimal development of the ovary. Thus, in 45,X and 46,XY females the ovaries are almost invariably devoid of oocytes. In contrast, germ cells in the testes do best when only one X chromosome is present; rarely do they survive in the XX or XXY condition.

When non-Y-bearing oocytes enter the differentiating gonad, the primary sex cords break up and encircle the oocytes in the cortex of the gonad (in contrast to the structure of the XY gonad). This occurs at approximately 16 weeks of gestation, and the isolated cell clusters are called primordial follicles. No new oogonia form after birth and many of them degenerate well before birth. Those that remain grow and become primary follicles to be stimulated following puberty.

2


UPPER GENITAL TRACT DEVELOPMENT

Phenotypic gender is determined by a complex tissue differentiation process that begins in the medial genital thickening or ridges on the posterior surface of the embryonic body cavity. Once gonadal sexual differentiation has begun, several other events must occur for normal male or female phenotypic differentiation to occur. During the fifth week after conception, coelomic epithelium, later known as germinal epithelium, thickens in the area of the medial aspect of the mesonephros. As germinal epithelial cells proliferate, they invade the underlying mesenchyme, producing the gonadal ridge. In the sixth week after conception the primordial germ cells, which formed at approximately the fourth week after conception, in the wall of the yolk sac, migrate up the dorsal mesentery of the hindgut and enter the undifferentiated gonad. These cells will differentiate into testes or ovaries based on the gene functions noted in [Chapter 1, Sexual Differentiation](#).

Signaled by the arrival of primordial germ cells in the fifth week after conception, two sets of paired genital ducts, the mesonephric or nephric (wolffian) ducts and the paramesonephric (müllerian) ducts, develop. The mesonephric system is the precursor to the male genital system and the paramesonephric to the female reproductive structures. The mesonephros is a prominent excretory structure that consists of a series of mesonephric tubules. The tubules connect with the elongating mesonephric (wolffian) ducts as the latter extend caudally, terminating in the urogenital sinus on each side of the midline. Derived from the evagination of the coelomic epithelium, the paramesonephric ducts develop lateral to each of the mesonephric ducts. The cephalward ends of these ducts open directly into the peritoneal cavity, whereas the distal ends grow caudally, fuse in the lower midline, and form the uterovaginal primordium. They join the urogenital sinus as an elevation, known

as the müllerian tubercle, which separates the urogenital area from the more posterior gut. Under the influence of the *SRY* gene in the prototestis the mesonephric (wolffian) ducts are maintained during development. As the developing male Sertoli cells begin to differentiate in response to *SRY*, they secrete a glycoprotein hormone, müllerian-inhibiting substance (MIS) or antimüllerian hormone (AMH), which causes the paramesonephric (müllerian) ducts to regress rapidly between the 8th and 10th fetal weeks. Without testosterone and AMH the mesonephric ducts degenerate and disappear, and the paramesonephric ducts develop into a uterus, fallopian tubes, and upper vagina. Leydig cells synthesize insulin-like-3 (coded by the *INSL3* gene) to promote transabdominal testicular descent into the scrotum. Mutations in this gene may lead to cryptorchidism. In females a structure similar to the gubernaculum develops in the inguinal canal, giving rise to the round ligaments that suspend the uterus in the adult.

Primary sex cords condense and extend to the medullary portion of the developing testes. They branch and join to form the rete testis. The testis therefore is primarily a medullary organ. Eventually the rete testis connects with the tubules of the mesonephric system and joins the developing epididymal duct. Müllerian duct remnants in the male include the appendix testis (hydatid of Morgagni) and the prostatic utricle. In females, MIS is not present, so müllerian ducts remain and the mesonephric tubules and ducts degenerate in the absence of androgens. This often results in remnant epoöphoron and paroöphoron cystic structures within the ovarian mesentery and Gartner duct cysts within the anterolateral vaginal wall. These structures are clinically important because they may develop into sizable and symptomatic cysts (see [Chapter 105, Vaginal Cysts](#)).



F. Netter M.D.

Figure 2.1 Homologs of the internal genitalia

The process of development and loss of the müllerian and wolffian systems begins at approximately the sixth week after conception and proceeds in a cephalad to caudal fashion. The more cephalad portions of the paramesonephric ducts, which open directly into the peritoneal cavity, form the fallopian tubes. The fused portion or uterovaginal primordium gives rise to the epithelium and glands of the uterus and cervix. Endometrial stroma and myometrium are derived from adjacent mesenchyme. Failure of the development of the paramesonephric ducts leads to agenesis of the cervix and uterus. Failure of fusion of the caudal portion of these ducts may lead to a variety of uterine anomalies, including complete duplication of the uterus and cervix or partial duplication of a variety of types (see Section VI, [Chapter 136](#), Uterine Anomalies:

Bicornuate, Septate, and Unicornuate Uterus). Peritoneal reflections in the area adjacent to the fusion of the two paramesonephric ducts give rise to the broad ligaments. Mesenchymal tissue here develops into the parametria.

The remnants of the mesonephric duct in the female include a small structure called the appendix vesiculosa, a few blind tubules in the broad ligaments (the epoöphoron), and a few blind tubules adjacent to the uterus (collectively called the paroöphoron). Remnants of the mesonephric duct system are often present in the broad ligaments and may be present adjacent to the uterus and/or vagina as Gartner duct cysts. The epoöphoron or paroöphoron may develop into cysts. Cysts of the epoöphoron are known as paraovarian cysts.




Figure 3.1 Homologs of external genitalia

imperforate until late in the embryonic life. However, occasionally, perforation does not completely occur (imperforate hymen). Failure of the sinovaginal bulbs to form leads to agenesis of the vagina. The precise boundary between the paramesonephric and urogenital sinus portions of the vagina has not been established.

Beginning in the fourth week after conception, the genital tubercle develops at the ventral tip of the cloacal membrane, with the labioscrotal swellings and urogenital folds developing soon after on either side of the cloacal membrane. In both sexes the genital tubercle subsequently elongates to form a phallus. By the end of the sixth week, the cloacal membrane is joined by the urorectal septum. This septum separates the cloaca into the urogenital sinus ventrally and the anal canal and rectum dorsally. The point on the cloacal membrane where the urorectal septum fuses will become the site of the perineal body. The cloacal membrane, now in two parts, then ruptures, opening the vulva and anal canal. Failure of the anal membrane to rupture results in an imperforate anus. With the opening of the urogenital membrane a urethral groove forms on the undersurface of the phallus, completing the undifferentiated portion of external genital development. Differences between male

and female embryos can be observed as early as the ninth week, but the distinct final forms are not found until 12 weeks of gestation.


Feminization of the undifferentiated external genitalia occurs in the absence of androgenic stimulation. The embryonic phallus remains quiescent and becomes the clitoris. The urogenital folds remain unfused except in front of the anus, forming the posterior fourchette. The unfused urogenital folds form the labia minora, while the labioscrotal folds remain as the labia majora. The labioscrotal folds fuse anteriorly to form the mons pubis. A portion of the urogenital sinus between the level of the hymen and the labia develops into the vestibule of the vagina, into which the urethra, the vagina, and the ducts of Bartholin glands enter. Beyond 12 weeks of gestation, the labioscrotal folds will no longer fuse when exposed to androgens, although other manifestations of masculinization may occur.

In contrast to the long-held belief that the development of the female genital was passive and occurred in the absence of androgens, the large number of estrogen receptors found in the genital tissues suggests that there is a role for maternal estrogens in the development of the female external genitalia. Female external

genital structures also contain androgen receptors. The distribution of androgen receptors resembles that of the male, which explains why the female genitalia can be masculinized if exposed to high androgen levels early in gestation.

The auxiliary genital glands of the female genitalia form from buds that grow out of the urethra. These buds come from the

surrounding mesenchyme and form the urethral and paraurethral glands (Skene glands). These glands correspond to the prostate gland in males. Similar outgrowths of the urogenital sinus form the vestibular glands (Bartholin glands), which are homologous to the bulbourethral glands in the male.



Tanner Stages of Breast Development




Figure 4.1 Developmental stages of the breast

Anatomy

- 5 External Genitalia
- 6 Perineum
- 7 Vagina
- 8 Pelvic Viscera
- 9 Cervix, Uterus, and Adnexa
- 10 Ovaries
- 11 Breast



This page intentionally left blank

The perineum is bound in front by the mons veneris, behind by the buttocks, and laterally by the thighs. More deeply it is limited by the margins of the pelvic outlet; namely, the pubic symphysis and arcuate ligament, ischiopubic rami, ischial tuberosities, sacrotuberous ligaments, sacrum, and coccyx. The vulva includes the portions of the female genital tract that are externally visible in the perineal region.

The mons veneris is a fatty prominence that overlies the symphysis pubis. It is covered by curly sexual (pubic) hair that functions as a dry lubricant during intercourse. From the mons veneris the labia majora extend in elliptical fashion to enclose the vulval cleft. They contain an abundance of adipose tissue and sebaceous and sweat glands and are covered by hair on their upper outer surfaces. Posteriorly a slightly raised connecting ridge, the posterior commissure or fourchette, joins them. Between the fourchette and the vaginal orifice a shallow, boat-shaped depression, the fossa

navicularis, is evident. The labia minora are thin, firm, pigmented, redundant folds of skin, which split anteriorly to enclose the clitoris. Laterally, they bound the vestibule and diminish gradually as they extend posteriorly. The skin of the small labia is devoid of hair follicles, poor in sweat glands, and rich in sebaceous glands.

The clitoris, a small, cylindrical, erectile organ situated at the lower border of the symphysis, is composed of two crura, a body and a glans. The crura lie deeply in close apposition with the periosteum of the ischiopubic rami. They join to form the body of the clitoris, which extends downward beneath a loose prepuce and is capped by the acorn-shaped glans. Generally, only the glans of the clitoris is externally visible between the two folds formed by the bifurcation of the labia minora.

The vestibule becomes apparent on separation of the labia. Within it are found the hymen, the vaginal orifice, the urethral meatus, and the opening of Skene and Bartholin ducts. The external

Pudendal, Pubic, and Inguinal Regions






Figure 5.1 External genitalia